

Repurposing a Cyclin-Dependent Kinase 1 (CDK1) Mitotic Regulatory Network to Complete Terminal Differentiation in Lens Fiber Cells

Allen Taylor,^{1,2} Yumei Gu,¹ Min-Lee Chang,¹ Wenxin Yang,¹ Sarah Francisco,¹ Sheldon Rowan,^{1,2} Eloy Bejarano,^{1,*} Steven Pruitt,^{3,*} Liang Zhu,⁴ Grant Weiss,⁵ Lisa Brennan,⁶ Marc Kantorow,⁶ and Elizabeth A. Whitcomb¹

¹Laboratory for Nutrition and Vision Research, USDA Human Nutrition Research Center on Aging, Tufts University, Boston, Massachusetts, United States

²Department of Ophthalmology, Tufts University School of Medicine, Boston, Massachusetts, United States

³Roswell Park Cancer Institute, Buffalo, New York, United States

⁴Albert Einstein College of Medicine, New York City, New York, United States

⁵Department of Neuroscience Tufts University School of Medicine, Boston, Massachusetts, United States

⁶Florida Atlantic University, Boca Raton, Florida, United States

Correspondence: Elizabeth A. Whitcomb, 711 Washington St., Boston, MA 02111, USA; elizabeth.whitcomb@tufts.edu. Allen Taylor, 711 Washington St., Boston, MA 02111, USA; allen.taylor@tufts.edu.

*Steven Pruitt deceased, June 11, 2019.

Current affiliation: **School of Health Sciences Universidad CEU Cardenal Herrera, Valencia, Spain.

Received: July 29, 2022

Accepted: November 22, 2022

Published: February 3, 2023

Citation: Taylor A, Gu Y, Chang ML, et al. Repurposing a cyclin-dependent kinase 1 (CDK1) mitotic regulatory network to complete terminal differentiation in lens fiber cells. *Invest Ophthalmol Vis Sci.* 2023;64(2):6. <https://doi.org/10.1167/iovs.64.2.6>

PURPOSE. During lens fiber cell differentiation, organelles are removed in an ordered manner to ensure lens clarity. A critical step in this process is removal of the cell nucleus, but the mechanisms by which this occurs are unclear. In this study, we investigate the role of a cyclin-dependent kinase 1 (CDK1) regulatory loop in controlling lens fiber cell denucleation (LFCD).

METHODS. We examined lens differentiation histologically in two different vertebrate models. An embryonic chick lens culture system was used to test the role of CDK1, cell division cycle 25 (CDC25), WEE1, and PP2A in LFCD. Additionally, we used three mouse models that express high levels of the CDK inhibitor p27 to test whether increased p27 levels affect LFCD.

RESULTS. Using chick lens organ cultures, small-molecule inhibitors of CDK1 and CDC25 inhibit LFCD, while inhibiting the CDK1 inhibitory kinase WEE1 potentiates LFCD. Additionally, treatment with an inhibitor of PP2A, which indirectly inhibits CDK1 activity, also increased LFCD. Three different mouse models that express increased levels of p27 through different mechanisms show impaired LFCD.

CONCLUSIONS. Here we define a conserved nonmitotic role for CDK1 and its upstream regulators in controlling LFCD. We find that CDK1 functionally interacts with WEE1, a nuclear kinase that inhibits CDK1 activity, and CDC25 activating phosphatases in cells where CDK1 activity must be exquisitely regulated to allow for LFCD. We also provide genetic evidence in multiple in vivo models that p27, a CDK1 inhibitor, inhibits lens growth and LFCD.

Keywords: CDK1, CDC25, WEE1, lens, differentiation, p27

During the life of an organism, tissues undergo cellular proliferation, differentiation, and aging. The differentiation process is particularly exaggerated in cells that discard their nucleus such as red blood cells, keratinocytes, and eye lens fiber cells. The lens of the eye must undergo this differentiation process to achieve optical clarity.

Within the lens, epithelial cells proliferate in an area anterior to the equator, where they differentiate into fiber cells. In this unidirectional process, lens fiber cells elongate, exit the cell cycle, and change their gene expression profile to primarily express crystallin proteins, similar to red blood cells synthesizing hemoglobin as their major gene product. As lens fiber cells mature, they compact in the center of the lens and lose their organelles in an ordered fashion.¹ The mechanisms by which each type of

organelle is removed are still being revealed. In support of roles for autophagy in removal of some organelles, it has been observed that organelle removal can be accelerated via enhancing autophagy by suppression of MAPK/JNK and PI3K signaling in a chick lens culture model.^{2,3} In the lens, removal of mitochondria, endoplasmic reticulum (ER), and Golgi requires the action of the mitophagy protein BNIP3L/NIX⁴ and lysosomal enzymes.⁵ However, the absolute requirement for distinct macroautophagy pathways in lens organelle removal remains unresolved as mouse models deleted for specific macroautophagy components exhibit no defect in organelle removal.^{6,7}

The last organelle to be removed during fiber cell differentiation is the nucleus. This process is initiated by disassembly of the nuclear membrane and completed with DNA




FIGURE 1. Regulation of CDK1 activity. Inhibitors are in red, and activators are shown in green. PP2A indirectly inhibits CDK1 by activating the inhibitory kinases WEE1 and MYT1 and inhibiting the CDK1 activating CDC25 phosphatases.

degradation by the lens-specific DNase, DLAD/DNase II β .⁸ The pathways that control nuclear envelope breakdown during lens fiber cell denucleation (LFCD) have not been fully elucidated. It has recently been proposed that LFCD in the chick involves a subcellular structure called the nuclear excisome.⁹ A number of cell cycle regulatory molecules are expressed in the nondividing lens fiber cells,^{10–13} and defects in cell division have been correlated with cataracts in humans,¹⁴ suggesting that mitotic cell cycle regulators may play a role in the unidirectional processes involved in establishing lens clarity. During the cell cycle, the activity of the essential CDK1 controls progression through mitosis.¹⁵ We hypothesized that vertebrate lenses utilize mitotic machinery, specifically CDK1, to direct phosphorylation of lamin and nuclear envelope breakdown during LFCD. Indeed, a lens-specific knockout confirmed a critical role for CDK1 in LFCD in the mouse.¹⁶

CDK1 activity in mitosis is regulated in multiple ways (Fig. 1). CDK1 activity is inhibited by phosphorylation at Tyr15 directed by the WEE1 and MYT1 kinases, while CDK1 activity is promoted by the CDC25 family of phosphatases, which remove the phosphorylation at Tyr15. The PP2A family of phosphatases regulates CDK1 activity indirectly by activating WEE1 and inhibiting the CDC25 family of phosphatases.¹⁷ These regulatory molecules have well-described roles in controlling the cell cycle and proliferation,^{18–20} but a role for their activity in lens differentiation has never been explored.

WEE1 is an essential kinase as the knockout is lethal early in development,²¹ and roles for WEE1 in cancer and mitosis have been well described.²⁰ Overexpression of WEE1 results in a “rough eye” phenotype in *Drosophila*,²² but a specific role for WEE1 in the vertebrate lens has not been described.

There are three members of the CDC25 family of phosphatases. While all three are known to regulate CDK function, only CDC25A is an essential gene and knockout embryos die at E5–E7,²³ suggesting CDC25 plays a role very early in development.²⁴ Intriguingly, CDC25B is regulated by the Alzheimer’s associated protein A β in primary human lens epithelial cells and is necessary for the increased resistance to oxidative stress stimulated by A β .²⁵ Furthermore, there is case report of an individual with a natural knockout of CDC25B who presented with a cataract.²⁶ Although there is no report of a lens phenotype in CDC25B knockout mice, these data suggest that CDC25B specifically may be important in maintenance of the lens.

PP2A is a family of phosphatases with multiple A, B, and C subunits and regulates multiple signaling pathways in

the cell. Knockout of either the C α or A α subunits is lethal between E6 and E10.²⁷ There is a report of knockdown of PP2A in developing goldfish impairing lens development,²⁸ suggesting that regulation of phosphorylation, potentially of CDK1, may be important in lens differentiation.

CDK1 activity is also inhibited by direct binding of the CDK inhibitor, p27(Kip1/CDKN1B).²⁹ p27 regulates cell cycle progression through its inhibition of CDK1.³⁰ Additionally, p27 is involved in CDK-dependent control of autophagy and apoptosis.³¹ However, the regulation of cell migration by p27 is independent of its ability to inhibit CDK.³² Increased levels of p27 were correlated with impaired LFCD in several mouse models of cataract.^{33,34}

Here we used mammalian and avian models to test the hypothesis that there is a dynamic CDK1 regulatory loop including positive regulation by the CDC25 family of phosphatases and negative regulation by WEE1, PP2A, and p27 that controls the nuclear disassembly and terminal differentiation process in lens fiber cells.

METHODS

Animals

All animals were bred and housed according to National Institutes of Health (NIH), Tufts University Institutional Animal Care and Use Committee (IACUC), and ARVO guidelines. Mice expressing a doxycycline-inducible p27 transgene were described previously.³⁵ Pregnant mice were treated for 2 or 5 days before birth to induce p27 expression during lens differentiation. Offspring were collected at day P1. Whole heads were fixed in 4% paraformaldehyde and subsequently dehydrated in sucrose before embedding in optimal temperature cutting compound (OCT). Heads of E18.5 embryos of wild-type (WT), *Skp2*^{-/-}, and p27197AA knock-in mice³⁶ were embedded in paraffin, sectioned, and stained with hematoxylin and eosin by the Tufts Medical Center histology core facility.

White Leghorn chicken eggs (Charles River Laboratories, Storrs, CT, USA) were sacrificed at days E8 to E18 in compliance with NIH, Tufts University IACUC, and ARVO guidelines. Primary chicken lens organ cultures were prepared from the lenses of embryonic day 12 as described in Brennan et al.³⁷ Briefly, whole lenses were removed from isolated chicken eyes and placed into individual wells of 24-well plates containing Medium 199 (ThermoFisher, Waltham, MA, USA) supplemented with 10% fetal bovine serum (Hyclone, Logan, UT, USA) and penicillin-streptomycin antibiotic mix (50 units/ml; ThermoFisher). Lenses were cultured for 2 hours prior to treatment to ensure only clear and undamaged lenses were used for culture studies. After 3 days of culture, lenses were collected and frozen in OCT.

Drugs used for chick lens culture included adavosertib (Selleckchem, Houston, TX, USA),³⁸ BN82002 (EMD Millipore, Burlington, MA, USA),³⁹ rubratoxin A (Cayman Chemical, Ann Arbor, MI, USA),⁴⁰ NSC663284 (Chem Cruz, Dallas, TX, USA),⁴¹ FTY720 (Chem Cruz),⁴² RO3306 (Sigma, St. Louis, MO, USA),⁴³ and CGP 745144⁴⁴ (Tocris Bioscience, Bristol, UK). Drugs were added at the beginning of culture and replenished daily.

Immunostaining

Chick lenses were fresh frozen in OCT prior to sectioning. Immediately prior to staining, sections were fixed

in 4% paraformaldehyde. Nuclei were visualized by Prolong mounting media (Invitrogen/ThermoFisher). In some cases, citrate-based antigen retrieval (Vector Laboratories, Burlingame, CA, USA) was necessary to elucidate staining. Lenses were sectioned serially, and thus the presence of multiple sections on a slide indicated that the intervening slides in the series (8 or 12) also had sections with areas of denucleation. Thus, the denucleation area was calculated by measuring the distance between nuclei, the number of sections with a denucleated region, the thickness of the section, and the number of slides in the series.

Antibodies

Antibodies included p27 mouse monoclonal (BD Biosciences, Franklin Lakes, NJ, USA), rabbit polyclonal (Santa Cruz Biotechnology, Dallas, TX, USA), phospholamin A (Abcam, Waltham, MA, USA), CDC25B (Santa Cruz Biotechnology), WEE1 (Santa Cruz Biotechnology), phospho-CDK1 (Y15; Cell Signaling Technologies, Danvers, MA, USA), DNase II β (kind gift from Shigekazu Nagata,

Osaka University), CDK1 (BD Biosciences), Ki67 (Vector Laboratories), CDC25B (Santa Cruz Biotechnology), CDC25C (Santa Cruz Biotechnology), and pCDC25C (Cell Signaling Technologies).

RESULTS

CDK1 as a Central Hub for Regulation of Lens Fiber Cell Denucleation

We previously demonstrated that CDK1 is necessary for efficient LFCD in the mammalian lens.¹⁶ To explore for the generalizability of this role for CDK1 in LFCD, we asked about development in an evolutionarily distant avian model. As in mice, chick lenses were surrounded on their anterior surface by a layer of epithelial cells (Fig. 2A, 2B). By embryonic day 8 (E8), mostly elongated nuclei were observed in the chick lens fibers at the center of the lens (Fig. 2A). By E10, more nuclei began to condense and became round in the central fiber cells (arrows). By day E14, all the nuclei in the center of the lens were condensed, and by E18,




FIGURE 2. Nuclear morphology and CDK1 expression in chick lens development. (A) DAPI staining of full section of E8 chick lens showing epithelium and fiber cell nuclei. E8 to E10 show central fibers; orange arrows show condensed nuclei in E10 and E12 sections. Scale bars are 50 μm. Panel for E18 represents a larger region than shown for other stages. (B) Chick lens section from a control lens (E12) placed in culture for 3 days with an organellar-free zone (OFZ) stained with CDK1 and DAPI; composite of fourteen 20x images. The OFZ appears offset due to section angle. Regions 3, 4, and 5 are ~785 μm, ~985 μm, and ~1100 μm from left edge of equatorial epithelium, respectively. Regions 1–5, 100x images: overlay and red channel (CDK1) shown as paired images. White line in region 5 is edge of OFZ. Arrows denote CDK1 expressing nuclei at edge of OFZ. Scale bars in 100x images are 20 μm. (C) Western blot of CDK1 and pCDK1 during chick lens development.




FIGURE 3. Inhibition of CDK1 activity impairs lens fiber cell denucleation. Chick lenses were put into culture at E12 for 3 days in the presence or absence of inhibitors as shown. DAPI staining of central fibers. Box-and-whisker plots: control (blue); drug treatment (orange). (x) median, line = mean. (A) Treatment with 20 μM RO-3306, $n = 10$. (B) treatment with 100 nM CGP 74514, $n = 5$. Full-size images of lens sections and data normalized to lens diameter are shown in Supplementary Fig. S1.

we observed a nuclei-free zone, indicating that LFCD had occurred, comparable to the sequence of development in the mammalian lens.

CDK1 was highly expressed in the cytoplasm and nuclei of chick lens anterior and equatorial epithelial cells (Fig. 2B, regions 1, 2). Furthermore, CDK1 was expressed in the nuclei of fiber cells through differentiation until just before denucleation (Fig. 2B, regions 3–5; see arrows), suggesting that CDK1 may have a role in LFCD in the chick as it does in the mouse.

During mitosis, the activity of CDK1 is tightly regulated by phosphorylation and binding to inhibitory proteins.⁴⁵ Elucidation of regulation of CDK1, particularly with regard to differentiation, has never been pursued in the chick lens. CDK1 activity is inhibited by phosphorylation at Tyr15. Western blot analysis indicates that during chick lens development, the levels of phosphorylated CDK1 decrease precipitously between E10 and E14 (Fig. 2C). This upregulation of CDK1 activity is consistent both with increased proliferation in the epithelium as well promoting LFCD in fiber cells.

We sought to confirm that CDK1 activity was required for efficient LFCD in chick lenses. Chick lenses continue their usual developmental program when maintained in vitro.^{2,3,37} Chick lenses were dissected at E12 and put in culture medium for 3 days in the presence or absence of CDK1 inhibitors. Inhibition of CDK1 with two different selective inhibitors, RO3306⁴³ and CGP 74514,⁴⁴ profoundly delayed LFCD (Figs. 3A, 3B). As gauged by region of denu-

cleation, inhibition of CDK1 delayed LFCD by four- to eight-fold (Fig. 3B, Supplementary Fig. S1). Treatment of lenses with inhibitors did not affect lens size as the diameter of the central sections was not affected by treatment (diameter of control for RO-3306, $1651 \pm 191 \mu\text{M}$; RO-3306, $1627.96 \pm 118 \mu\text{M}$; diameter of control for CGP, $74,514 = 1547 \pm 121 \mu\text{M}$; CGP, $74,514 = 1586 \pm 157 \mu\text{M}$), suggesting that treatment affects denucleation rather than growth. These inhibitors showed activity toward CDK1 in cultured cells as we observed increased phosphorylated CDK1 upon treatment with RO3306 and CGP 74514 (Supplementary Fig. S2, lanes 1, 2, 6, 7). Together, these findings indicate that as in the mouse, CDK1 controls LFCD in the chick.¹⁶

A Regulatory Loop of CDC25, PP2A, and WEE1 Regulates CDK1 and LFCD

CDK1 phosphorylation, and thus activity, is controlled by a network of kinases and phosphatases (Fig. 1). Thus, we asked if these regulators were also involved in LFCD. The family of CDC25 phosphatases removes the Tyr15 phosphorylation on CDK1 and promotes its activity.¹⁷ Immunofluorescence analysis indicates that CDC25B is colocalized with CDK1 in nuclei just outside the organelle-free zone in chick lenses (Fig. 4A), suggesting that CDC25B regulates the activity of CDK1 in these cells. Additional support for a role for CDC25B in LFCD comes from the observation




FIGURE 4. CDC25 regulates lens fiber cell denucleation. (A) Immunofluorescence of chick lens showing CDC25B (green) and CDK1 (red) 20 \times and 100 \times . (B, C) E12 chick lenses placed in culture for 3 days in the presence or absence of inhibitors as shown. (B) 10 μ M NSC663284, $n = 4$. (C) 10 μ M BN82002, $n = 4$. Full-size images of lens sections and data normalized to lens diameter are shown in Supplementary Fig. S4.

that CDC25B is colocalized with the nuclear pore complex in central lens fiber cells, and they disappear coordinately before denucleation (Supplementary Fig. S3A). Furthermore, CDK1 and CDC25B physically interact, as demonstrated by coimmunoprecipitation with α CDK1 (Supplementary Fig. S3B). CDC25B is also expressed in fiber cell nuclei prior to LFCD in mouse lenses (Supplementary Fig. S3C) suggesting a common pathway of CDK1 regulation in the mouse and chick.

If LFCD requires CDC25, then inhibition of this phosphatase should inhibit LFCD. To determine the role of CDC25 in LFCD, we treated chick lenses with the selective CDC25 inhibitors NSC663284⁴¹ and BN82002.³⁹ As expected, cells treated with these drugs show increased levels of p-CDK1, indicating inhibition of CDK1 activity through inhibition of CDC25 (Supplementary Fig. S1A,

lanes 1, 3, 6, 8). Treatment with inhibitors did not affect lens growth as the diameter of the central sections was not affected (diameter of control for NSC663284, $1499 \pm 245 \mu$ m; NSC663284, $1587 \pm 146.76 \mu$ m; diameter of control for BN82002, $1514 \pm 117 \mu$ m; BN82002, $1658 \pm 118 \mu$ m). Fiber cells in lenses treated with NSC663284 failed to denucleate and BN82002 delayed LFCD markedly (Figs. 4B, 4C, Supplementary Fig. S4). Combined, the data indicate that CDK1 activation by CDC25 is necessary for efficient LFCD.

WEE1 kinase phosphorylates CDK1 on Tyr15, thereby inhibiting its activity.¹⁷ Immunofluorescence analysis indicates that WEE1 expression overlaps with CDC25B expression (Fig. 5A), suggesting that like CDC25, WEE1 is part of the regulatory machinery of CDK1 and of LFCD. In contrast with inhibition of CDK1 and CDC25 leading to inhibition of LFCD, it was anticipated that inhibition of WEE1 would




FIGURE 5. Inhibition of WEE1 potentiates lens fiber cell denudation. (A) Immunofluorescence of chick lens showing localization of CDC25B (green) and WEE1 (red). Arrows show coordinate disappearance. Images are 20 \times and 100 \times . (B) E12 lenses were placed in culture in the presence and absence of 1 μ M adavosertib as noted, $n = 6$. Mean line for control (blue) is at the bottom of the box. Full-size images of lens sections and data normalized to lens diameter are shown in Supplementary Fig. S5.

activate CDK1 and accelerate LFCD. In support of this hypothesis, treatment of lenses with adavosertib accelerated LFCD by 60% (Fig. 5B, Supplementary Fig. S5) but did not affect overall lens growth (diameter of control, $1696 \pm 26 \mu\text{M}$; adavosertib, $1649 \pm 115 \mu\text{M}$) indicating an inhibitory role for WEE1 in LFCD. Confirming its activity with regard to WEE1 regulation of CDK1, adavosertib decreased the level of p-CDK1 in cells in culture (Supplementary Fig. S2, lanes 4, 5).

PP2A is a multisubunit phosphatase that activates WEE1 and MYT1 while inhibiting CDC25 (Fig. 1); thus, we hypothesized that activation of PP2A would inhibit CDK1 and delay LFCD, whereas inhibition of PP2A would accelerate LFCD. To test this hypothesis, lenses were treated with a PP2A activator FTY720 and inhibitor rubratoxin A. As predicted, the PP2A inhibitor caused a 60% increase in LFCD (Fig. 6A, Supplementary Fig. S6). In comparison, activation of PP2A decreased LFCD 40% (Fig. 6B, Supplementary Fig. S6). Treatment of cells in culture with the PP2A activator increased p-CDK1 while the PP2A inhibitor, rubratoxin, decreased p-CDC25⁴⁰ (Supplementary Fig. S2, lanes 6, 9–11). Treatment of lenses with either rubratoxin A or FTY720 did not affect lens growth (diameter of control for rubratoxin A, $1708 \pm 154 \mu\text{M}$; rubratoxin A, $1577 \pm 131.74 \mu\text{M}$; diameter of control for FTY720, $1699 \pm 76 \mu\text{M}$; FTY720, $1622.7 \pm 133 \mu\text{M}$).

Collectively, these results indicate that CDK1 in fiber cell nuclei interacts with an inhibiting kinase, WEE1, and an activating phosphatase, CDC25, in the region where CDK1 activity is exquisitely regulated to foster LFCD. The activating phosphatase and inhibiting kinase are themselves regulated by PP2A, and thus LFCD is regulated by PP2A activity. Furthermore, these data support a fine-tuned spatiotemporal regulation of CDK1 with positive and negative regulators of CDK1 expressed in the lens fiber cells immediately prior to denudation.

p27 Regulates CDK1 in Postproliferative Developing Fibers and Inhibits LFCD

In mitosis, p27 inhibits CDK1 by direct binding and thus functions in a regulatory capacity different from WEE1 and CDC25. Increased levels of p27 have been correlated with impaired LFCD in several mouse models of cataract, but direct evidence for a mechanistic connection between p27 and LFCD is lacking.^{33,34} Thus, we hypothesized that increased p27 would directly interfere with LFCD. We tested this hypothesis using three different mouse models.

To begin, we determined whether transgenic overexpression of p27 interferes with efficient LFCD using an inducible




FIGURE 6. Inhibition of PP2A potentiates lens fiber cell denucleation, while activation of PP2A inhibits denucleation. (A, B) E12 chick lenses were placed into culture in the presence and absence of drugs for 3 days as noted. (A) 2 μ M rubratoxin, $n = 7$. (B) 2.5 μ M FTY720, $n = 5$. Full-size images of lens sections and data normalized to lens diameter are shown in Supplementary Fig. S6.

p27 transgene. Mice were treated with doxycycline at E15.5 or E18.5 of gestation, and offspring were examined at P1. Mice not exposed to doxycycline showed a clear organelle-free zone upon birth (Fig. 7A, left panels). In comparison, induction of p27 for 2 days from E18.5 to birth showed increased p27 levels and retained nuclei in the putative organelle-free zone (Fig. 7A, right panels). These findings indicate that increased expression of p27 impairs denucleation. Lenses in mice that were treated for 5 days (from day E15.5 to birth) showed even more p27 expression and also exhibited impaired LFCD (Supplementary Fig. S7A). Lenses from mice expressing p27 for 2 days, were $\sim 35\%$ smaller than WT lenses, and lenses from mice expressing p27 for 5 days were $\sim 60\%$ smaller. Confirming a p27-CDK1 block of cell proliferation lenses from these mice decreased staining of the proliferation marker Ki67 (Fig. 7B, 1Supplementary Fig. S7B).

An anticipated consequence of inhibition of CDK1 by increased levels of p27 is decreased phosphorylation of lamin A associated with retained nuclear membranes and impaired LFCD. As predicted, overexpression of p27 decreased the level of phosphorylated lamin A in the lens (Fig. 7C, upper panels). This was associated with retention of nuclei (Fig. 7C, lower panels). Staining of p27 was observed in the control epithelium (arrows in 7C) and in fiber cells when fluorescence intensity was increased (Supplementary Fig. S7C).

Degradation of DNA, among the final steps in LFCD, is accomplished by a lens-specific enzyme, DLAD/DNAse II β ,⁸ that digests genomic material following nuclear envelope

breakdown. In control mice, DLAD/DNAse II β was allowed into the nucleus in the center of the lens (Fig. 7D, left panel, inset 2) but excluded from peripheral nuclei (Fig. 7D, left panel, inset 1). Consistent with retained nuclear membranes in lenses with elevated p27, DLAD/DNAse II β was excluded from central nuclei from p27 overexpressing mice as it was in outer zones of WT mice (Fig. 7D, right panel, see insets 1, 2).^{16,33,34} This recapitulates observations when CDK1 is absent.¹⁶

p27 levels are regulated in part by ubiquitination with subsequent degradation by the proteasome. The ubiquitin ligase complex SCF^{Skp2} utilizes a substrate binding subunit called the F-box to ubiquitinate a number of cell cycle regulatory molecules, including p27.^{46–49} We examined the lenses from embryos from which the F-box substrate targeting subunit *Skp2* was deleted. As expected, lenses from *Skp2*^{-/-} showed increased levels of p27 (Supplementary Fig. S8). Lenses from *Skp2*^{-/-} embryos were $\sim 50\%$ smaller than the WT lenses (Fig. 8 compare A1 to B1), consistent with a regulatory role in epithelial cell proliferation and the decreased growth phenotype of the mice.³⁶ Nevertheless, the overall lens structure was preserved. There was a single layer of epithelia, a discernable bow region and a posterior region free of epithelial cells. The equatorial (region 2) and anterior (region 3) epithelium regions appeared normal in these knockout mice. However, *Skp2*^{-/-} embryos showed retained cell nuclei in the presumptive LFCD organelle-free zone (region 4). These findings indicate that *Skp2* deletion has a major regulatory effect on LFCD but perhaps a limited effect on overall lens formation.




FIGURE 7. Expression of an inducible p27 transgene impairs lens fiber cell denucleation. Doxycycline-responsive p27 transgene was induced during gestation for 0, 2, or 5 days. Offspring were analyzed at P1. Le, lens epithelium; lc, lens cortex. **(A)** Lens sections were analyzed for p27 (red, upper panels) and DNA/DAPI (lower panels). **(B)** Staining of Ki67⁺ cells in lens epithelium in control (left) and 2-day induction (right). **(C)** P1 lenses from control (left) or 2-day p27 overexpressor mice (right) were stained for phospho-lamin A (green), p27 (red), and DNA/DAPI as indicated. Arrowheads show p27⁺ nuclei in control epithelium. Lower panels show DNA/DAPI staining in the center of the lens, broken line in panel above. **(D)** P1 lenses from control (left) and p27 two-day overexpressor mice (right) were stained for DLAD/DNase IIβ. Insets show peripheral (1) and central (2) nuclei.

A requirement for SCF^{Skp2}-catalyzed ubiquitination and degradation of p27 is phosphorylation at Thr187 of p27. We also examined lenses from embryos with a homozygous p27^{T187A} mutation that precludes T187 phosphorylation (T187AA⁵⁰). Similar to *Skp2*^{-/-}, lenses from embryos expressing this degradation incompetent mutant p27 showed increased levels of p27 (Supplementary Fig. S8), and nuclei were retained in the center of the lens (Fig. 8C, region 4). Additionally, the lenses were ~50% smaller than the WT lenses. As with the *Skp2*^{-/-} lenses, the equatorial (Fig. 8C, region 2) and anterior epithelium (region 3) of the T187AA lenses appeared morphologically normal, indicating that *Skp2*-dependent targeting of p27 does not affect the overall development of the lens. These results indicate that inhibiting the degradation of p27 inhibits the activity of the essential regulatory molecule CDK1 and impairs LFCD.

DISCUSSION

The function of the lens is to focus light on the retina. Lens clarity is established and maintained by the expression and organization of high levels of crystallin proteins in the lens fibers along with the programmed removal of organelles in the central region. Failure to remove organelles is correlated with cataracts in animal models as well as in humans.^{5,14,34,51} Previous work indicated that the mitotic kinase CDK1 has been adapted to initiate LFCD.¹⁶ Here we asked if other processes that affect CDK1 activity in mitosis are also involved in LFCD in evolutionarily distant vertebrates. We demonstrate that a mitotic network including p27, WEE1, CDC25, and PP2A regulates CDK1 activity in the lens and tightly controls the process of LFCD in both the mouse and chick. This research establishes new biological roles for these widely expressed regulatory enzymes.




FIGURE 8. Impaired degradation of p27 leads to inefficient denucleation. E18.5 embryos from (A) WT, (B) *Skp2*^{-/-}, and (C) p27 T187AA knock-in mice were examined by hematoxylin and eosin. (1) Whole lens, (2) equatorial epithelium, (3) anterior epithelium, and (4) central fibers. Three individual embryos of each genotype were analyzed with similar results.

First, we observed that as in the mouse, CDK1 activity is upregulated in chick lens development. Furthermore, CDK1 was expressed in fiber cell nuclei in the chick until denucleation, and two inhibitors of CDK activity impaired LFCD in embryonic chick lenses. These data indicate that CDK1 activity is a master regulator of LFCD in both the mouse and chick.

CDK1 activity was controlled by hierarchical and counterbalanced phosphorylation and dephosphorylation activities of WEE1/MYT1 and CDC25 (Fig. 1). The colocalization of CDK1 with inhibitor WEE1 and activator CDC25B in nuclei just before nuclear envelope breakdown suggested that the activity of CDK1 was tightly regulated by these molecules in nuclei just before LFCD. We corroborated this pharmacologically: treatment of lens with drugs that inhibited the CDC25 family of phosphatases impaired LFCD, indicating that CDC25 activity is essential for activation of CDK1 during LFCD. While the inhibitors were active against all family members, the colocalization of CDC25B with CDK1 and prior literature indicating a role for CDC25B specifically in lens function suggest that CDC25B is the relevant phosphatase in LFCD.^{25,26} It would be of interest to determine if there is a denucleation defect in the *CDC25B*^{-/-} animals, although there are no reports of ocular defects. Inhibition of WEE1 activity with adavosertib potentiated LFCD, indicating that CDK1 activity is suppressed by WEE1 during lens development.

In mitotic pathways, the activities of WEE1 and CDC25 are mechanistically linked via the PP2A phosphatase, which inhibits CDC25 and activates WEE1. As expected, inhibition of PP2A with rubratoxin A potentiated LFCD, while activation of PP2A with FTY720 inhibited LFCD. Although PP2A inhibits CDK1 activation and thus nuclear envelope breakdown, we cannot rule out whether other pathways regulated by PP2A are also contributing to the defect in LFCD we observe after FTY720 treatment.

In addition to regulating LFCD, CDK1 regulated lens epithelial cell division, which might also affect lens size.¹⁶ However, we did not observe a difference in lens size between treatment and control during the lens culture. This

may not be unexpected as we did not observe alterations in lens size in the lens-specific *CDK1* knockout mice.¹⁶ Changes in the lens mass during the culture period are primarily due to postmitotic fiber cell differentiation and increases in cell size due to the significant upregulation of crystallin expression⁵² and both intrinsic and extrinsic factors,⁵³ including compaction.⁵⁴ We cannot rule out whether inhibition of CDK1 activity in the epithelium is indirectly affecting fiber cell differentiation, and this would be interesting to explore.

Previously, by perturbing the ubiquitin proteasome system by expression of a mutant ubiquitin in the lens, we observed increased levels of the CDK inhibitor p27 correlated with impaired LFCD, and this was corroborated in multiple mouse models of cataract.^{33,34} Here in addition to directly showing that increased levels of p27 impair LFCD, we demonstrate that retarding degradation of p27 also results in decreased LFCD. These data indicate that increased p27 levels directly inhibit LFCD. In confirmation of these findings, DLAD/DNase II β , which usually enters the disassembled nuclei to degrade chromatin, is excluded from nuclei in lenses with increased levels of p27. Together, these observations suggest a CDK1-regulated LFCD process comparable to that which regulates the cell cycle (Fig. 1).

While organelle removal including the mitochondria, ER, and nucleus can be accelerated by stimulating autophagy,^{2,3} mice deficient in autophagy proteins are not defective in lens organelle removal^{6,7}; thus, the role of autophagy in these processes remains to be explored further. Elimination of ER, Golgi, and mitochondria can be separated from nuclear removal as mouse models with deletions of BNIP3L/NIX or SNFH2 show impaired removal of one organelle but not the other.^{4,55} Our data indicate that mitotic regulators are controlling nuclear removal during lens differentiation, and we did not observe any alterations in the removal of ER and mitochondria upon CDK1 knockout, further confirming removal of these organelles are controlled by different pathways.¹⁶

Evidence indicates that lysosomal activities are required for organelle removal. Lysosomes are a source of DLAD/DNase II β to digest DNA that remains after

the nuclear membrane breakdown. Additionally, lysosomal phospholipases are critical for organelle removal in zebrafish and mouse models.⁵ Whereas deletion of *Plaati1* in zebrafish impaired the degradation of nonnuclear organelles, deletion of the homologous phospholipase *PLAAT3* in the mouse impaired removal of organelles, including nuclei. The authors suggest that the activity of *Plaati1/PLAAT3* is involved in lysosomal membrane breakdown, and the lack of sufficient DLAD/DNAse II β might be the cause of impaired LFCd in these mice.⁵ However, initiation of LFCd requires that the nuclear membrane be permeabilized prior to access of nuclear material to DLAD/DNAse II β . Thus, there must be upstream regulators of nuclear membrane disassembly. We postulate that phosphorylation of nuclear lamins by CDK1 is responsible for permeabilization of the nuclear membrane, perhaps priming it for further destruction by phospholipases.

Nonmitotic nuclear envelope breakdown has been observed in other tissues. Keratinocytes and erythroid progenitor cells both remove their nuclei during differentiation.⁵⁶ However, the mechanisms by which they remove nuclei are distinct, relying on different kinases to direct nuclear disassembly, and thus they are examples of convergent evolution. Neutrophils use the process of NETosis, during which part of the nucleus breaks off and is extruded from the cell, as a means to destroy pathogens and to remodel vasculature.^{57,58} NETosis is dependent on CDK4/6 and is regulated by the CDK inhibitor p21.⁵⁹ Furthermore, there is disassembly of cytoskeletal networks and remodeling of nuclear lamins in NETosis as in mitosis and LFCd.⁵⁸ Thus, our work provides further elucidation of nonmitotic nuclear envelope breakdown involving a mitotic regulatory network.

Acknowledgments

The authors thank Jonas Galper for use of his laboratory for chick experiments and Giuliana Perini-Villanueva for expert technical assistance.

Supported by NIH RO1EY028559 and RO1EY026979 (to A.T.) and USDA 8050-51000-089-01S (A.T.). This material is based upon work supported by the US Department of Agriculture–Agricultural Research Service (ARS), under Agreement Nos. 58-1950-4-003 and 58-8050-9-004.

Disclosure: **A. Taylor**, None; **Y. Gu**, None; **M.-L. Chang**, None; **W. Yang**, None; **S. Francisco**, None; **S. Rowan**, None; **E. Bejarano**, None; **S. Pruitt**, None; **L. Zhu**, None; **G. Weiss**, None; **L. Brennan**, None; **M. Kantorow**, None; **E.A. Whitcomb**, None

References

1. Bassnett S. On the mechanism of organelle degradation in the vertebrate lens. *Exp Eye Res.* 2009;88:133–139.
2. Basu S, Rajakaruna S, Reyes B, Van Bockstaele E, Menko AS. Suppression of MAPK/JNK-MTORC1 signaling leads to premature loss of organelles and nuclei by autophagy during terminal differentiation of lens fiber cells. *Autophagy.* 2014;10(7):1193–1211.
3. Gheyas R, Ortega-Alvarez R, Chauss D, Kantorow M, Menko AS. Suppression of PI3K signaling is linked to autophagy activation and the spatiotemporal induction of the lens organelle free zone. *Exp Cell Res.* 2022;412(2):113043.
4. Brennan LA, McGreal-Estrada R, Logan CM, Cvekl A, Menko AS, Kantorow M. BNIP3L/NIX is required for elimination of mitochondria, endoplasmic reticulum and Golgi apparatus during eye lens organelle-free zone formation. *Exp Eye Res.* 2018;174:173–184.
5. Morishita H, Eguchi T, Tsukamoto S, et al. Organelle degradation in the lens by PLAAT phospholipases. *Nature.* 2021;592(7855):634–638.
6. Morishita H, Eguchi S, Kimura H, et al. Deletion of autophagy-related 5 (Atg5) and Pik3c3 genes in the lens causes cataract independent of programmed organelle degradation. *J Biol Chem.* 2013;288(16):11436–11447.
7. Matsui M, Yamamoto A, Kuma A, Ohsumi Y, Mizushima N. Organelle degradation during the lens and erythroid differentiation is independent of autophagy. *Biochem Biophys Res Commun.* 2006;339(2):485–489.
8. Nishimoto S, Kawane K, Watanabe-Fukunaga R, et al. Nuclear cataract caused by a lack of DNA degradation in the mouse eye lens. *Nature.* 2003;424(6952):1071–1074.
9. Costello MJ, Brennan LA, Mohamed A, Gilliland KO, Johnsen S, Kantorow M. Identification and ultrastructural characterization of a novel nuclear degradation complex in differentiating lens fiber cells. *PLoS One.* 2016;11(8):e0160785.
10. Gao CY, Bassnett S, Zelenka PS. Cyclin B, p34cdc2, and H1-kinase activity in terminally differentiating lens fiber cells. *Dev Biol.* 1995;169(1):185–194.
11. Gao CY, Rampalli AM, Cai HC, He HY, Zelenka PS. Changes in cyclin dependent kinase expression and activity accompanying lens fiber cell differentiation. *Exp. Eye Res.* 1999;69(6):695–703.
12. He HY, Gao C, Vrensen G, Zelenka P. Transient activation of cyclin B/Cdc2 during terminal differentiation of lens fiber cells. *Dev Dyn.* 1998;211(1):26–34.
13. Zhang P, Wong C, DePinho RA, Harper JW, Elledge SJ. Cooperation between the Cdk inhibitors p27(KIP1) and p57(KIP2) in the control of tissue growth and development. *Genes Dev.* 1998;12(20):3162–3167.
14. Gulluni F, Prever L, Li H, et al. PI(3,4)P2-mediated cytokinetic abscission prevents early senescence and cataract formation. *Science.* 2021;374(6573):eabk0410.
15. Santamaria D, Barriere C, Cerqueira A, et al. Cdk1 is sufficient to drive the mammalian cell cycle. *Nature.* 2007;448(7155):811–815.
16. Chaffee BR, Shang F, Chang ML, et al. Nuclear removal during terminal lens fiber cell differentiation requires CDK1 activity: Appropriating mitosis-related nuclear disassembly. *Development.* 2014;141(17):3388–3398.
17. Potapova TA, Daum JR, Pittman BD, et al. The reversibility of mitotic exit in vertebrate cells. *Nature.* 2006;440(7086):954–958.
18. Brenner AK, Reikvam H, Lavecchia A, Bruserud O. Therapeutic targeting the cell division cycle 25 (CDC25) phosphatases in human acute myeloid leukemia—the possibility to target several kinases through inhibition of the various CDC25 isoforms. *Molecules.* 2014;19(11):18414–18447.
19. Crncec A, Hochegger H. Triggering mitosis. *FEBS Lett.* 2019;593(20):2868–2888.
20. Ghelli Luserna di Rora A, Cerchione C, Martinelli G, Simonetti G. A WEE1 family business: Regulation of mitosis, cancer progression, and therapeutic target. *J Hematol Oncol.* 2020;13(1):126.
21. Tominaga Y, Li C, Wang RH, Deng CX. Murine Wee1 plays a critical role in cell cycle regulation and pre-implantation stages of embryonic development. *Int J Biol Sci.* 2006;2(4):161–170.
22. Price DM, Jin Z, Rabinovitch S, Campbell SD. Ectopic expression of the Drosophila Cdk1 inhibitory kinases, Wee1 and Myt1, interferes with the second mitotic wave and disrupts pattern formation during eye development. *Genetics.* 2002;161(2):721–731.

23. Ray D, Terao Y, Nimbalkar D, et al. Hemizygous disruption of Cdc25A inhibits cellular transformation and mammary tumorigenesis in mice. *Cancer Res.* 2007;67(14):6605–6611.
24. Bouldin CM, Kimelman D. Cdc25 and the importance of G2 control: Insights from developmental biology. *Cell Cycle.* 2014;13(14):2165–2171.
25. Xu J, Li D, Lu Y, Zheng TY. Abeta monomers protect lens epithelial cells against oxidative stress by upregulating CDC25B. *Free Radic Biol Med.* 2021;175:161–170.
26. Lam CW, Fong NC, Chan TY, et al. Centrosome-associated CDC25B is a novel disease-causing gene for a syndrome with cataracts, dilated cardiomyopathy, and multiple endocrinopathies. *Clin Chim Acta.* 2020;504:81–87.
27. Reynhout S, Janssens V. Physiologic functions of PP2A: Lessons from genetically modified mice. *Biochim Biophys Acta Mol Cell Res.* 2019;1866(1):31–50.
28. Liu WB, Hu XH, Zhang XW, et al. Protein serine/threonine phosphatase-2A is differentially expressed and regulates eye development in vertebrates. *Curr Mol Med.* 2013;13(8):1376–1384.
29. Lacy ER, Filippov I, Lewis WS, et al. p27 binds cyclin-CDK complexes through a sequential mechanism involving binding-induced protein folding. *Nat Struct Mol Biol.* 2004;11(4):358–364.
30. Yoon MK, Mitrea DM, Ou L, Kriwacki RW. Cell cycle regulation by the intrinsically disordered proteins p21 and p27. *Biochem Soc Trans.* 2012;40(5):981–988.
31. McKay LK, White JP. The AMPK/p27(Kip1) pathway as a novel target to promote autophagy and resilience in aged cells. *Cells.* 2021;10(6):1430.
32. Rampioni Vinciguerra GL, Citron F, Segatto I, Belletti B, Vecchione A, Baldassarre G. p27kip1 at the crossroad between actin and microtubule dynamics. *Cell Div.* 2019;14:2.
33. Caceres A, Shang F, Wawrousek E, et al. Perturbing the ubiquitin pathway reveals how mitosis is hijacked to denude and regulate cell proliferation and differentiation in vivo. *PLoS One.* 2010;5:e13331.
34. Lyu L, Whitcomb EA, Jiang S, et al. Unfolded-protein response-associated stabilization of p27(Cdkn1b) interferes with lens fiber cell denudation, leading to cataract. *FASEB J.* 2016;30(3):1087–1095.
35. Pruitt SC, Freeland A, Rusiniak ME, Kunnev D, Cady GK. Cdkn1b overexpression in adult mice alters the balance between genome and tissue ageing. *Nat Commun.* 2013;4:2626.
36. Zhu L, Skoultchi AI. Coordinating cell proliferation and differentiation. *Curr Opin Genet Dev.* 2001;11(1):91–97.
37. Brennan L, Disatham J, Kantorow M. Hypoxia regulates the degradation of non-nuclear organelles during lens differentiation through activation of HIF1a. *Exp Eye Res.* 2020;198:108129.
38. Hirai H, Iwasawa Y, Okada M, et al. Small-molecule inhibition of Wee1 kinase by MK-1775 selectively sensitizes p53-deficient tumor cells to DNA-damaging agents. *Mol Cancer Ther.* 2009;8(11):2992–3000.
39. Brezak MC, Quaranta M, Mondesert O, et al. A novel synthetic inhibitor of CDC25 phosphatases: BN82002. *Cancer Res.* 2004;64(9):3320–3325.
40. Wada S, Usami I, Umezawa Y, et al. Rubratoxin A specifically and potentially inhibits protein phosphatase 2A and suppresses cancer metastasis. *Cancer Sci.* 2010;101(3):743–750.
41. Lazo JS, Aslan DC, Southwick EC, et al. Discovery and biological evaluation of a new family of potent inhibitors of the dual specificity protein phosphatase Cdc25. *J Med Chem.* 2001;44(24):4042–4049.
42. Nagahara Y, Matsuoka Y, Saito K, Ikekita M, Higuchi S, Shinomiya T. Coordinate involvement of cell cycle arrest and apoptosis strengthen the effect of FTY720. *Jpn J Cancer Res.* 2001;92(6):680–687.
43. Vassilev LT, Tovar C, Chen S, et al. Selective small-molecule inhibitor reveals critical mitotic functions of human CDK1. *Proc Natl Acad Sci USA.* 2006;103(28):10660–10665.
44. Dai Y, Dent P, Grant S. Induction of apoptosis in human leukemia cells by the CDK1 inhibitor CGP74514A. *Cell Cycle.* 2002;1(2):143–152.
45. Domingo-Sananes MR, Kapuy O, Hunt T, Novak B. Switches and latches: A biochemical tug-of-war between the kinases and phosphatases that control mitosis. *Philos Trans R Soc Lond B Biol Sci.* 2011;366(1584):3584–3594.
46. Carrano AC, Eytan E, Hershko A, Pagano M. SKP2 is required for ubiquitin-mediated degradation of the CDK inhibitor p27. *Nat Cell Biol.* 1999;1(4):193–199.
47. Hara T, Kamura T, Nakayama K, Oshikawa K, Hatakeyama S. Degradation of p27(Kip1) at the G(0)-G(1) transition mediated by a Skp2-independent ubiquitination pathway. *J Biol Chem.* 2001;276(52):48937–48943.
48. Nakayama K, Nagahama H, Minamishima YA, et al. Targeted disruption of Skp2 results in accumulation of cyclin E and p27(Kip1), polyploidy and centrosome overduplication. *Embo J.* 2000;19(9):2069–2081.
49. Nakayama K, Nagahama H, Minamishima YA, et al. Skp2-mediated degradation of p27 regulates progression into mitosis. *Dev Cell.* 2004;6(5):661–672.
50. Malek NP, Sundberg H, McGrew S, et al. A mouse knock-in model exposes sequential proteolytic pathways that regulate p27Kip1 in G1 and S phase. *Nature.* 2001;413(6853):323–327.
51. Boniuk M, Zimmerman LE. Ocular pathology in the rubella syndrome. *Arch Ophthalmol.* 1967;77(4):455–473.
52. Bloemendal H, de Jong W, Jaenicke R, Lubsen NH, Slingsby C, Tardieu A. Ageing and vision: Structure, stability and function of lens crystallins. *Prog Biophys Mol Biol.* 2004;86(3):407–485.
53. Bassnett S, Sikic H. The lens growth process. *Prog Retin Eye Res.* 2017;60:181–200.
54. Bassnett S, Costello MJ. The cause and consequence of fiber cell compaction in the vertebrate lens. *Exp Eye Res.* 2017;156:50–57.
55. He S, Limi S, McGreal RS, et al. Chromatin remodeling enzyme Snf2h regulates embryonic lens differentiation and denudation. *Development.* 2016;143(11):1937–1947.
56. Rogerson C, Bergamaschi D, O'Shaughnessy RFL. Uncovering mechanisms of nuclear degradation in keratinocytes: A paradigm for nuclear degradation in other tissues. *Nucleus.* 2018;9(1):56–64.
57. Binet F, Cagnone G, Crespo-Garcia S, et al. Neutrophil extracellular traps target senescent vasculature for tissue remodeling in retinopathy. *Science.* 2020;369(6506):eaay5356.
58. Thiam HR, Wong SL, Qiu R, et al. NETosis proceeds by cytoskeleton and endomembrane disassembly and PAD4-mediated chromatin decondensation and nuclear envelope rupture. *Proc Natl Acad Sci USA.* 2020;117(13):7326–7337.
59. Amulic B, Knackstedt SL, Abu Abed U, et al. Cell-cycle proteins control production of neutrophil extracellular traps. *Dev Cell.* 2017;43(4):449–462.e5.