

## SYSTEMATIC ARTICLE

# Herbal Extracts with Antifungal Activity against *Candida albicans*: A Systematic Review

Hsuan Hsu<sup>1</sup>, Chirag C. Sheth<sup>2</sup> and Veronica Veses<sup>3,\*</sup>

<sup>1</sup>Department of Dentistry, Faculty of Health Sciences, Universidad Cardenal Herrera, CEU Universities, Moncada 46113, Valencia, Spain; <sup>2</sup>Department of Medicine, Faculty of Health Sciences, Universidad Cardenal Herrera, CEU Universities, Moncada 46113, Valencia, Spain; <sup>3</sup>Department of Biomedical Sciences, Faculty of Health Sciences, Universidad Cardenal Herrera, CEU Universities, Moncada 46113, Valencia, Spain

**Abstract:** In the era of antimicrobial resistance, fungal pathogens are not an exception. Several strategies, including antimicrobial stewardship programs and high throughput screening of new drugs, are being implemented. Several recent studies have demonstrated the effectiveness of plant compounds with antifungal activity. In this systematic review, we examine the use of natural compounds as a possible avenue to fight fungal infections produced by *Candida albicans*, the most common human fungal pathogen. Electronic literature searches were conducted through PubMed/MEDLINE, Cochrane, and Science Direct limited to the 5 years. A total of 131 articles were included, with 186 plants extracts evaluated. Although the majority of the natural extracts exhibited antifungal activities against *C. albicans* (both *in vivo* and *in vitro*), the strongest antifungal activity was obtained from *Lawsonia inermis*, *Pelargonium graveolens*, *Camellia sinensis*, *Mentha piperita*, and *Citrus latifolia*. The main components with proven antifungal activities were phenolic compounds such as gallic acid, thymol, and flavonoids (especially catechin), polyphenols such as tannins, terpenoids and saponins. The incorporation of nanotechnology greatly enhances the antifungal properties of these natural compounds. Further research is needed to fully characterize the composition of all herbal extracts with antifungal activity as well as the mechanisms of action of the active compounds.

---

## ARTICLE HISTORY

---

Received: February 20, 2020  
Revised: May 19, 2020  
Accepted: May 21, 2020

DOI:  
10.2174/1389557520666200628032116

**Keywords:** Herbal extracts, antifungal properties, *Candida albicans*, gallic acid, thymol, catechin, nanotechnology.

## 1. INTRODUCTION

*Candida* species are commensal microorganisms in human oral mucosa, digestive and vaginal tracts. Normally, people with healthy immune systems can control the growth and spread of this opportunistic fungus. However, when the host becomes weak and immunocompromised, it can lead to serious infection. These infections can be superficial such as thrush, vaginitis, skin infections, or invade the bloodstream and spread to any site of the human host, which can cause many clinical complications such as brain abscess, endocarditis, meningitis, arthritis and pyelonephritis [1].

Approximately 150 *Candida* species have been identified, of which only about 20 species can cause infection in humans. Among these, *Candida albicans* is the most common pathogenic species responsible for most invasive infections in immunocompromised patients, followed by *Candida glabrata*,

*Candida tropicalis*, *Candida parapsilosis*, and *Candida krusei*, which comprised up to 90% of *Candida* infections [2]. It is estimated that more than a quarter of a million patients are infected with invasive candidiasis, with the incidences rates up to 2-14 per 100000 populations globally [3]. In addition, *Candida* is ranked fourth of the most common pathogens of bloodstream infections after *Staphylococcus aureus* and *Enterococci* [4]. Furthermore, *C. albicans* is one of the most isolated species responsible for nosocomial infections due to the use of intravenous catheters, invasive procedures, transplantation, wide range use of broad-spectrum antibiotics and chemotherapies [2]. Particular characteristics of *C. albicans* are their morphological transition between yeast and hyphal forms, which allow adherence to oral mucosa; formation of biofilms; phenotypic switching and the secretion of virulence factors such as adhesins and hydrolytic enzymes [5].

Currently, there are only five major classes of antifungal agents available to treat *C. albicans* infections. These include polyenes, allylamines and azoles, which target ergosterol and nucleoside analogues, which inhibit DNA and/or RNA syn-

---

\*Address correspondence to this author at the Department of Biomedical Sciences, Faculty of Health Sciences, Universidad Cardenal Herrera, CEU Universities, Moncada 46113, Valencia, Spain; Tel: +3496136900 ext 64351; E-mail: [veronica.veses@uchceu.es](mailto:veronica.veses@uchceu.es)

thesis, and echinocandins, that inhibit the synthesis of  $\beta$ -1, 3-glucan [6]. Among these, azole antifungals such as fluconazole are often selected as the treatment choice because they are well tolerated, exhibit low toxicity, available for oral administration, and are inexpensive. However, in recent years, the resistance of *Candida* species to antifungal drugs has increased worldwide [7]. Generally, antifungal resistance is achieved through reduced intracellular drug accumulation, decreased target affinity for the drug and counteraction of the drug effect.

Due to the widespread and overuse of limited antifungal drugs, the search for alternatives against *C. albicans* is ongoing, especially in plants and natural herbs. It is estimated that there are 250,000-500,000 species of plants on earth and only 10% are used by humans [8], which provided 50% of commercially available modern drugs [9]. Plant extract therapies have been utilized and accepted all over the world due to their low side effects. The earliest record of plant medications can be traced back to 2600 BCE in Mesopotamia, Egypt, India, Greece and China, revealing about 300-1000 different drugs [9]. The most common species used by ancient people are algae, bryophytes, pteridophytes and angiosperms [9]. Different geographical locations also have a big impact on the development of herbal drug systems and the availability of plant resources. Kier and coworkers describe that the highest diversity of plants may be found in the Neotropic (central and south America) and the Asia-Pacific region (China, India, USA, Australia), and lower diversities in Africa and on oceanic islands [10]. Traditionally, the majority of medicinal plants are found in India and China, while Europe and the USA have developed fewer sources [9]. The diversity of plants provides a wide range of important sources of biologically active molecules with enormous potential antifungal properties, such as phenols, tannins, terpenoids and alkaloids [9]. Isolated and modified compounds such as dimethyl pyrrole, hydroxydihydrocornin-aglycones and indole derivatives have also shown antifungal activity *in vitro* [8]. Studies have reported that the extraction method of active substances has a great influence on the function of antimicrobial components and their antifungal effectiveness. Silver nanoparticles, antibodies, and photodynamic inactivation have increased the distribution and effectiveness of antifungal drugs [2]. For this reason, the antifungal activity of natural herbs and extracts have been assessed as an alternative antifungal drug against *C. albicans*. In this systematic review, we evaluate the antifungal activities of natural herbs and extracts and their synergistic effect with common antifungal agents.

## 2. MATERIALS AND METHODS

### 2.1. Search Strategy

This review was carried out in accordance with PRISMA guidelines. Comprehensive, structured literature searches were conducted *via* the databases PubMed/MEDLINE, Cochrane and Science Direct. The publication date was limited from Jan 1<sup>st</sup>, 2015 until Feb 23<sup>rd</sup>, 2019. The electronic search was performed using the phrases: *C. albicans* AND (extract OR herbal OR natural) AND (antifungal) for Science Direct and Cochrane Library; Search terms with Mesh

terms: (*C. albicans*) AND extracts [MeSH Terms]) OR natural products [MeSH Terms]) OR herbal [MeSH Terms]) AND antifungal [MeSH Terms]) were searched in PubMed in the English language.

### 2.2. Inclusion Criteria

The fundamental inclusion factors were that the studies must involve the use of natural products, herbs, or extract against *C. albicans*. Studies could be either *in vitro*, *in vivo*, or both for the purpose of assessing the antifungal activity of natural products against *C. albicans*. Table 1 shows all the study inclusion and exclusion criteria.

### 2.3. Types of Study

All prospective or longitudinal studies, experimental studies, clinical trial/study, double-blinded, randomized, placebo-controlled trials examining nature were included.

### 2.4. Types of Preparation

Herbal preparations are described as naturally prepared from herbs or plants from their roots, flowers, leaves, fruits, bulbs or seeds through different extraction methods into essential oil or extracts. These were then applied to inactive placebo or active control such as common antifungal drugs (azoles, nystatin, amphotericin B). Studies combining herbal interventions and routine pharmacologic therapy (co-intervention) were also reviewed.

### 2.5 Selection Criteria

Studies omitted from this review include retrospective studies, editorials, letters, reviews, case reports, cohort studies and pilot studies. Studies not using *C. albicans* as a tested organism and not presenting minimum inhibitory concentration values (MIC) were also excluded. Essential oils and extracts originating from animals or insects were excluded from this study.

### 2.6. Study Selection

Firstly, primary literature research was conducted. Next, the abstracts and titles were evaluated in order to screen and eliminate articles unrelated to this research topic. Following this, the remaining studies were downloaded as full-text articles and were assessed for eligibility. Only studies meeting the inclusion and exclusion criteria were included in this systematic review.

### 2.7. Data Extraction

Tables 2 and 3 were used to organize the information gained from each study [11-19]. Table 2 displays data from combined *in vivo/in vitro* studies whilst Table 3 contains information from *in vitro* studies only [20-141]. The following data was collected from all eligible articles: the scientific and common names of the plants; country of collection; parts of the plants that were extracted; extraction methods; strains that were tested; MIC or colony-forming units (CFU) of the products and outcomes.

Table 1. List of inclusion and exclusion criteria.

Inclusion Criteria	Exclusion Criteria
Study type: prospective or longitudinal studies, experimental studies, Clinical Trial, Clinical study, RCT (Only articles with level of evidence of 1b are included)	Retrospective study, editorials, letters, review, case report, cohort, pilot study
<i>C. albicans</i> strains must be tested	Algae/Animal/Insects as interventions.
Results must include antifungal assessment/evaluation method using MIC <i>in vitro</i> and <i>in vivo</i> . CFU unit is also included in <i>in vivo</i> studies.	Studies in which <i>C. albicans</i> are not tested.
Full length article available	-
English-language only	-
Published between Jan 1,2015 to Feb 23rd, 2019	-
Studies combined natural products with common antifungal drugs are included (Nystatin and azoles)	-

Table 2. Data extraction table from combination *in vivo/in vitro* studies investigating the antifungal activity of plant extracts against *C. albicans*.

Refs.	Plant/Organism (Common Name)	Country of Origin	Plant Part(s)	Product (s)	<i>In vivo</i> : MIC /CFU (mg/mL, mg/mL)	<i>In vitro</i> : MIC (mg/mL, mg/mL)	Host Organism	Strains	Conclusion
[11]	<i>Lawsonia inermis</i>	Iran	Leaves	EE	5-10 mg/mL	-	Wistar rats	<i>C. albicans</i> LC201976	4% was more effective than 2% and was as effective as clotrimazole.
[12]	<i>Punica granatum L.</i>	Algeria	-	AC	80 mg/mL	0.090 mg/mL	Male mice	<i>C. albicans</i> <i>C. krusei</i> <i>C. guilliermondii</i>	<i>Quercus suber L.</i> show the best and <i>Vicia faba</i> had the poor antifungal activity.
	<i>Quercus suber L.</i>				20 mg/mL	0.105 mg/mL			
	<i>Vicia faba</i>				>100 mg/mL	0.010mg/mL			
[13]	<i>Camellia sinensis</i> (L.) O Kuntze	Algeria	-	AC	40 mg/mL	5 µg /mL	C57BL6 mice	<i>C. albicans</i> , <i>C. glabrata</i> , <i>C. tropicalis</i> , <i>C. krusei</i>	AC was more active
				AQ	60 mg/mL	20 µg /mL			
[14]	<i>Morinda tomentosa</i>	Indonesia	Roots	ME	>32mg/mL	-	<i>Galleria mellonella</i>	<i>C. albicans</i> DSY2521 <i>C. albicans</i> CAF2-1	x
[15]	<i>Melaleuca alternifolia</i>	Brazil	-	Essential oil	5.33 Log <sub>10</sub> CFU	1.95 mg/mL	Male mice	<i>C. albicans</i> strain ATCC 18804	<ul style="list-style-type: none"> <li>12.5% extract concentration completely inhibited the biofilms</li> <li>Protective effect against oral <i>C. albicans</i> infections in mice.</li> </ul>
[16]	<i>Mitracarpus frigidus</i>	Brazil	Aerial	ME	400 and 4000mg kg <sup>-1</sup> CFU: Log 4.68 (day one)	500 µg /mL	Female Wistar rats	<i>C. albicans</i> ATCC 10231	<ul style="list-style-type: none"> <li>Promising antifungal activity <i>in vitro</i> and <i>in vivo</i>.</li> <li><i>In vitro</i> results suggest its ability to act on the cellular envelope.</li> <li>Better than fluconazole (MIC value = 10,000µg /mL)</li> </ul>

(Table 2) Contd....

Refs.	Plant/Organism (Common Name)	Country of Origin	Plant Part(s)	Product (s)	In vivo: MIC /CFU (mg/mL, mg/mL)	In vitro: MIC (mg/mL, mg/mL)	Host Organism	Strains	Conclusion
[17]	<i>Syzygium cumini</i>	Brazil	Seeds	NaP	100mg/kg	-	Diabetic infected Wistar rats	-	<ul style="list-style-type: none"> <li>Nanotechnology improve antioxidant properties.</li> <li>Contain high concentrations of phenols and flavonoids (gallic acid, chlorogenic acid, grutin, quercetin)</li> <li>Hypoglycaemic activity in rat models of DM</li> </ul>
[18]	<i>Astragalus membranaceus</i>	China	Roots	Low molecular weight polysaccharide (LMW-ASP)	CFU: 5.87 ± 0.03c -6.05 log10	-	Sera of mice infected with live <i>C. albicans</i> cells	-	Greatly improved against systemic candidiasis by strongly enhancing Th1 and Th2 responses in recombinant protein rP-HSP90C, but mechanism is unclear.
[19]	<i>Jatropha curcas L</i>	Mauritius	Barks, roots, leaves, seeds	Crude extracts	350-3290 mg/L	17.80-83.30 mg/mL	<i>Bactrocera zonata</i> and <i>B. cucurbitae</i> (Diptera fruit flies)	<i>C. albicans</i> ATCC 1023	<ul style="list-style-type: none"> <li>Show antifungal activity <i>in vivo</i> and <i>in vitro</i>.</li> <li>ME of mature leaves show the lowest activity and bark ME extract was highest <i>in vivo</i>.</li> <li>Contains alkaloid, steroids, tannins, flavonoids, phenol and coumarins.</li> </ul>

Abbreviations: \*-: Not specified/Not available, \*X: No antifungal effect; \*ME: Methanolic extract, \*AC: Acetone extract, \*AQ: Aqueous extract, \*AQE: Aqueous ethanolic extract; \*EtOAc: Ethyl acetate extract, \*EE: Ethanol extract, \*Dichloromethane extract: DCM, \*HE: Hexane extract; \*CHL: Chloroform extract, \*BA: butanol extract; \*NaP: Nanoparticles/Nano formulations.

## 2.8. Antifungal Activity Measurement

Both *in vitro* and *in vivo* antifungal activities are measured by Minimum Inhibitory Concentration (MIC), which is defined as the lowest concentration of an antimicrobial drug that will inhibit the visible growth of a microorganism after overnight incubation. Antifungal activities are measured either by microdilution assay, tube diffusion method or serial microplate dilution methods. In addition, *in vivo* studies are also assessed by quantifying the CFU, which is a measure used to estimate the number of viable bacteria or fungal cells in a sample.

## 2.9. Data Quality Evaluation

The quality of studies was evaluated according to the Centre for Evidence-based Medicine Levels of Evidence and PRISMA guidelines [149,150].

## 3. RESULTS AND DISCUSSION

### 3.1. Description of Selected Reports

Fig. (1) depicts an overview of the study selection procedure. After the removal of duplicates, a total of 2666 articles were recovered from three databases, with publication dates ranging from January 1, 2015, to February 23<sup>rd</sup>, 2019. Fol-

lowing the screening of titles and abstracts, 2283 articles were excluded, leaving 379 full-text articles, which were assessed for eligibility. Finally, a total of 131 articles met the inclusion criteria and were considered suitable for this systematic review. In total, there were 186 natural products involved in this systematic review; please see Tables 2 and 3 for further details on each study, [11-19] [20-141]. Plants identified in the review originated in 42 countries, with the largest percentages in Brazil (20%), India (9%) and Iran (7%) (Fig. 2). This geographical distribution and preference are supported by a study by Kier and coworkers [10].

Most herbal extracts show minimal to moderate antifungal effects against *C. albicans*; however, 27 tested plants were ineffective against it. In terms of herbal interventions, seven studies utilized nanotechnology with herbal extracts; ten articles assessed the synergistic effect of natural products with common antifungal drugs. Fourteen plants have been tested repeatedly in several studies and appear more than once in this review, which are *Salvadora persica*, *Camellia sinensis*, *Cinnamomum verum*, *Cuminum cyminum*, *Lawsonia inermis*, *Melaleuca alternifolia*, *Mentha piperita*, *Origanum vulgare*, *Paeonia lactiflora*, *Pelargonium graveolens*, *Psidium guajava*, *Syzygium aromaticum* and *Thymus vulgaris*.

**Table 3. Overview of names, countries of origin, plant part(s), formulation, MIC, strains used and conclusions of the herbal interventions *in vitro*.**

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, µg/mL)	Used Strains	Conclusion
[20]	<i>Olea europaea</i> (Olive)	Croatia	Leaves	Extract	46.875 mg/mL	<i>C. albicans</i> ATCC 10231 <i>C. dubliniensis</i> CBS 7987	<ul style="list-style-type: none"> <li>• Cytotoxic effect on tested yeast strains</li> <li>• Concentration dependent.</li> <li>• Contains hydroxytyrosol, protocatechuic acid, tyrosol, oleuropein, pinoresinol and apigenin.</li> </ul>
[21]	<i>Melaleuca alternifolia</i>	Australia	Leaves	Tea tree oil (TTO)	Average 0.19% Fluconazole +TTO: 38.46mg/mL	<i>C. albicans</i> ATCC 10231 <i>C. albicans</i> strains resistant to fluconazole	Show antifungal and synergistic effect with fluconazole
[22]	<i>Stryphnodendron adstringens</i> (Mart.) Coville (Leguminosae)	Brazil	Stem barks	Dried and pulverized	15.6 µg/mL	<i>C. albicans</i> ATCC 10231	Successfully inhibited planktonic growth and biofilm development.
[23]	<i>Metasequoia glyptostroboides</i>	Korea	Cone (abietane-type diterpenoid taxodone)	EtOAc	250-1000(mg/mL)	<i>C. albicans</i> KBN06P00076 <i>C. Albicans</i> KBN06P00074	Effective against <i>C. albicans</i>
[24]	<i>Eugenia dysenterica</i> DC. ( <i>Hexachlamys macedoi</i> Legrand) <i>Pouteria ramiflora</i> (Mart.) Radlk, <i>Pouteria torta</i> (Mart.) Radlk, <i>Bauhinia rufa</i> (Bong.) Steud, <i>Erythroxylum subrotundum</i> A	Brazil	Leaves	AQ	Cannot be detected	<i>C. guilliermondii</i> ATCC 6260, <i>C. tropicalis</i> ATCC 28707 <i>C. parapsilosis</i> ATCC 22019 <i>C. albicans</i> ATCC 90028, <i>C. Glabrata</i> ATCC 2001, <i>C. Famata</i> ATCC 62894 <i>C. krusei</i> ATCC 34135	<ul style="list-style-type: none"> <li>• No inhibition detected against <i>C. Albicans</i> and <i>C. Glabrata</i>.</li> <li>• AQ show significant inhibitory activity against <i>C. Parapsilosis</i>, <i>C. Guilliermondii</i>, <i>C. Tropicalis</i>, <i>C. Krusei</i> and <i>C. Famat</i>.</li> </ul>
[25]	<i>Piper guineense</i>	Nigeria	Fruits and leaves	AQ EE ME CHL HE	AQ: NA EE:78 µg/mL ME:39 µg/mL CHL:78 µg/mL HE:78 µg/mL	<i>C. albicans</i> ATCC 10231 <i>C. Glabrata</i> ATCC 2001 <i>C. Tropicalis</i> ATCC 750 <i>C. Parapsilosis</i> ATCC 7330	ME, EE, CHL and HE show antifungal efficacy, whereas AQ is not effective.
[26]	<i>Pelargonium graveolens</i>	USA	Purchased	Geranium oil (GO) Nanoemulsion geranium oil (NGE)	GO:1.82 µg/mL NGE:3.64 µg/mL	<i>C. albicans</i> ATCC 14053 <i>C. Tropicalis</i> ATCC 66029 <i>C. Glabrata</i> ATCC 66032 <i>C. Krusei</i> ATCC 6258	<ul style="list-style-type: none"> <li>• NGE was twice higher than the GO and more effective to reduce the amount of biofilm in the catheter.</li> <li>• Eliminate biofilm formation.</li> </ul>
[27]	<i>Swartzia simplex</i>	Panama	Root and bark	DCM	32 µg/mL	<i>C. albicans</i> DSY2621 Parent wild-type CAF2-1	Show antifungal activity
[28]	<i>Bursera morelensis</i>	Mexico	Stems	Ramirez essential Oil	0.062 – 0.25 mg/mL	<i>C. albicans</i> ATCC 14065 <i>C. Albicans</i> ATCC 32354	Germ tube inhibition and diminish the transcription of the gene <i>INT1</i> .

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, mg/mL)	Used Strains	Conclusion
[29]	<i>Aeollanthus cucullatus</i>	Cameroon		Hexane ethyl acetate extract	0.625 - 5 mg/mL	<i>C. albicans</i> <i>C. glabrata</i> .	Biofilm inhibition by blocking the filamentation process and by reducing the biofilm thickness.
[30]	<i>Leguminosae</i> family species	Brazil	Leaves	ME	-	<i>C. albicans</i> <i>C. krusei</i> <i>C. glabrata</i> <i>C. tropicalis</i> <i>C. parapsilosis</i> ,	None of them show antifungal activity.
[31]	<i>Ricinus communis</i>	Ghana	leaves	AQ ME EE	3.13 - 25.0 mg/mL	<i>C. albicans</i> .	<ul style="list-style-type: none"> <li>All are effective against <i>C. albicans</i>,</li> <li>ME show higher antifungal activity than other extract. may be due to the presence of high amount of tannins, flavonoids, and terpenoids.</li> </ul>
[32]	<i>Cochlospermum regium</i>	Brazil	Pilger roots	EtOAc	250 mg/mL	<i>C. albicans</i> 10231 <i>C. krusei</i> 34135 <i>C. glabrata</i> 2001 <i>C. tropicalis</i> 28707	Effective due to the presence of tannins and gallic acid.
[33]	<i>Salvia adenophora Fernald (Lamiaceae)</i>	Italy	Aerial	Isolated compounds	-	<i>C. albicans</i> clinical strain.	X
[34]	<i>Olea africana</i>	South Africa	Leaves	HE CHL DCM, EtOAc EE ME BA AQ	Average 0.37 mg/mL	<i>C. albicans</i>	All are effective however <i>C. neoformans</i> and <i>E. faecalis</i> were the most sensitive test organisms.
[35]	<i>Helichrysum</i> species: <i>H. armenium</i> DC, <i>H. arenarium</i> L. (Moench)	Turkey	-	EE	All are 8 µg /mL	<i>C. albicans</i> <i>C. parapsilosis</i>	<i>H. arenarium</i> is the most remarkable among other tested extracts.
[36]	<i>Antidesma madagascariense</i> Lam. ( <i>Euphorbiaceae</i> )	Mauritius	Leaves	AQ AC	4.00 mg/mL	<i>C. albicans</i> ATCC 10231	Show antifungal activity and show antioxidant, anti-inflammatory activity and serve as AChE inhibitors.
[37]	<i>Lavandula binaludensis</i> <i>Cuminum cyminum</i>	Iran	Aerial parts	Essential oils	7.91 mg/ mL  8.00 mg/mL	<i>C. albicans</i> isolates	<ul style="list-style-type: none"> <li>Effective against <i>C. albicans</i>.</li> <li>Antifungal are attributed to g-terpinene and 1,8-cineole through destroying cell walls and proteins, interfering in the work of membrane enzymes and affecting DNA and RNA replication.</li> </ul>
[38]	<i>Lawsonia inermis</i> <i>Withania somnifera</i> <i>onga</i> <i>Euphorbia hirta</i> <i>Pogostemon parviflorus</i> <i>Adenocalymma alliacum</i> ,	India	Leaves	ME EE	ME:2.8 mg/mL ME: 3.2 mg/mL EE: 3.1 mg/mL ME:5.0 mg/mL EE: 2.81 mg/mL ME:1.5 mg/mL EE: 2.75 mg/mL ME: 4.3 mg/mL EE: 4.25 mg/mL ME: 3.15 mg/mL EE: 3.85 mg/mL	<i>C. albicans</i>	<i>W. somnifera</i> , <i>C. longa</i> , <i>Euphorbia hirta</i> , <i>Echinophora platyloba</i> , <i>Zingiber officinale</i> , <i>L. inermis</i> , <i>Adenocalymma alliacum</i> , <i>P. parviflorus</i> and <i>Swertia chirata</i> effective against <i>C. albicans</i> at MIC 5 mg/mL without any toxic effect.

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, µg/mL)	Used Strains	Conclusion
	<i>Echinophora platybola</i>				EE: 2.3 mg/mL		
	<i>Zingiber officinale</i>				ME: 2.10 mg/mL EE: 4.25 mg/mL		
39]	<i>Thymus capitatus</i>	Tunisia	Aerial parts	Essential oils	125 µg/mL	<i>C. albicans</i> 328, <i>C. albicans</i> 247, <i>C. albicans</i> 311, <i>C. albicans</i> 249, <i>C. riferii</i> 648, <i>C. tropicalis</i> , <i>C. glabrata</i> 239, <i>C. glabrata</i> 113	<ul style="list-style-type: none"> <li><i>C. verum</i> exhibited the best activity; <i>S aromaticum</i> showed moderate activity; <i>P. graveolens</i> EO was less active.</li> <li>Affect ergosterol biosynthesis and disturb fatty acid homeostasis.</li> </ul>
	<i>Pelargonium graveolens</i>				250-1000 µg/mL		
	<i>Cinnamomum verum</i>				31.25-62.5 µg/mL		
	<i>Syzygium aromaticum</i>				125-250 µg/mL		
[40]	<i>Sideroxylon obtusifolium</i>	Brazil	leaves	Hydro alcoholic extracts	62.5 mg/mL	<i>C. albicans</i> ATCC 10231	Show antifungal activity in the presence of flavonoid and saponins.
	<i>Syzygium cumini</i>				125 mg/mL		
[41]	<i>Polyscias fulva (Hiern)</i>	Cameroon	Stem bark	Crude DCM ME	12.5 µg/mL 50 µg/mL 100 µg/mL	<i>C. albicans</i> ATCC 1663 <i>C. Krusei</i> ATCC 6258 <i>C. Parapsilosis</i> ATCC 22019 <i>C. Lucitaniae</i> ATCC 200950 <i>C. Glabrata</i> IP 35 <i>C. Guilliermondii</i> clinical isolate	Show antifungal activity due to the presence of terpenoid and saponins
[42]	<i>Ficus drupacea</i>	Egypt	Stem bark	7 isolated compounds and	7-7521 µg/mL	<i>C. albicans</i> ATCC 26555	<ul style="list-style-type: none"> <li>Isolate compounds show better antifungal activity than extract.</li> <li>Compounds 5-O-methylatifolin) and 7 (epilupeol acetate) exhibited the highest antifungal activity.</li> </ul>
				n-Hexane extract	4-15 µg/mL		
[43]	<i>Tamarix gallica</i>	Tunisia	Leaves and flowers	ME	0.292 mg/mL	<i>C. parapsilosis</i> , ATCC 22019 <i>C. Albicans</i> , ATCC 90026; <i>C. Krusei</i> ATCC 6258; <i>C. Glabrata</i> ATCC 90030	<ul style="list-style-type: none"> <li>Show antifungal activity.</li> <li>Flower presence flavonoids and leaves showed quercetin 3-O-glucuronide these suggest antifungal activity.</li> </ul>
44]	<i>Carissa opaca</i>	Pakistan	Root	ME Ethyl Acetate (EA)	ME: 20 mg/mL EA: 7.8 mg/mL	<i>C. albicans</i> ATCC 10231	ME showed moderate to high antimicrobial activities and EA displayed especially notable efficacy.
[45]	<i>Helichrysum microphyllum</i> subsp. <i>Tyrrhenicum</i>	Italy	-	Essential oils (ESO)	750 µg/mL	<i>C. albicans</i> ATCC 10231	<ul style="list-style-type: none"> <li>Chitosan formulations increased antifungal activity against <i>C. albicans</i>.</li> <li>Terpenes and alcohols such as c-curcumene and lina- examined is responsible for the antifungal effect</li> </ul>
				ESO with Chitosan	94.5 µg/mL		
[46]	<i>Rhaphiodon echinus</i> (Nees and Mart)	Brazil	Leaves	Essential oils	>1024 µg/mL	<i>C. albicans</i> , <i>C. Krusei</i> <i>C. Tropicalis</i>	X

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, mg/mL)	Used Strains	Conclusion
[47]	<i>Carya illinoensis</i>	Brazil	Leaves	AE EE	AE: 0.78 - 25 mg/mL EE: 1.56 - 25 mg/mL	<i>C. tropicalis</i> ATCC 66029 <i>C. parapsilosis</i> ATCC 22019 <i>C. albicans</i> ATCC 14053	<ul style="list-style-type: none"> <li>• AE and EE were effective at all concentrations by inactivating germ tube production.</li> <li>• Presence of phenolics acids (gallic acid and ellagic acid), flavonoids (rutin) and condensed tannin (catechin and epicatechin).</li> </ul>
[48]	<i>Buchenavia tetraphylla</i>	Brazil	Leaves	HE CHL EE ME	HE 156 - 2500 mg/mL CHL: 156-1250 mg/mL. ME: 625-1250 mg/mL EE :625-2500 mg/mL	<i>C. albicans</i> strains F01, F02,F03,F08, F11,F14,F22,F23,F27,UFPEDA 1007	ME showed the best activity which inhibit cell division and able enhance the action of fluconazole
[49]	<i>Corymbia intermedia</i>	Australia	Stem and leaves	AQE	500 µg/mL,	<i>C. albicans</i>	<ul style="list-style-type: none"> <li>• 80% AQEt of <i>S. glomulifera</i> was the most active.</li> <li>• The leaves of <i>S. glomulifera</i> contain antibacterial components: α-pinene, aromadendrene and globulol, eucalyptin, and compounds betulinic acid, oleanolic acid-3-acetate and ursolic acid-3-acetate</li> </ul>
	125 µg/mL						
	31 µg/mL						
[50]	<i>Ocimum basilicum</i>	Italy	-	Essential oils	0.09 - 4.58 mg/mL	<i>C. albicans</i> <i>C. famata</i>	<ul style="list-style-type: none"> <li>• <i>T. vulgaris</i> and <i>O. vulgare</i> essential oils showed the best activity against all the tested pathogens.</li> <li>• Rich in monoterpenes, carvacrol and thymol, there together can completely block ergosterol synthesis and making porous the membrane</li> <li>• <i>S. sclarea</i> showed no antifungal effect</li> </ul>
	<i>Origanum vulgare</i>				0.018 - 3.6 mg/mL		
	<i>Salvia sclarea</i>				No activity		
	<i>Thymus vulgaris</i>				0.09 – 1.87 mg/mL		
	<i>Illicium verum</i>				0.19 - >19.5 mg/mL		
[51]	<i>Plumbago rosea</i>	India	-	Plumbagin	5 µg/mL	<i>C. albicans</i> ATCC2091, <i>C. tropicalis</i> clinical isolate <i>C. parapsilosis</i> clinical isolate	Show antifungal activity by disrupting biofilm.
[52]	<i>Scabiosa arenaria</i>	Tunisia	Flowers, fruits, stems, leaves and roots	BA	0.02 mg/mL	<i>C. ATCC reference strains</i> , <i>C. albicans</i> ATCC 90028, <i>C. glabrata</i> ATCC 90030, <i>C. krusei</i> ATCC 6258, <i>C. parapsilosis</i> ATCC 22019.	<ul style="list-style-type: none"> <li>• Show antifungal activity.</li> <li>• Present of oleanolic acid and luteolin-7-O-glucoside show good antimicrobial effect.</li> </ul>
[53]	<i>Bersama abyssinica</i> , <i>Embelia schimperi</i> , <i>Ocimum lamiifolium</i> , <i>R. steudneri</i> <i>R. nepalensis</i> <i>Z. scabra</i>	Ethiopia	Leaves and roots	EE	512 mg/mL 512 mg/mL 512 mg/mL 512 mg/mL 512 mg/mL 512 mg/mL	<i>C. albicans</i>	<ul style="list-style-type: none"> <li>• 74% of the medicinal plant extracts tested exhibited antimicrobial effect against more of the 12 different microbial strains.</li> <li>• <i>E. schimperi</i>, <i>O. lamiifolium</i>, and <i>R. steudneri</i> was found to be the most promising plants against microbes.</li> </ul>

(Table 3) Contd....



Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, µg/mL)	Used Strains	Conclusion
[54]	<i>Euphorbia paralias L</i>	Tunisia	Stems and leaves	CHL	0.015 – 5 mg/mL	<i>C. albicans</i> ATCC 90028	Could be great potential as new antimicrobial agents.
[55]	<i>Eucalyptus globulus</i>	Brazil	Purchased	Essential oil (EO) Nanoemulsion (NE)	0.219 mg/mL 0.7 mg/mL	<i>C. albicans</i> <i>C. tropicalis</i> <i>C. glabrata</i>	<ul style="list-style-type: none"> <li>Nanoemulsion was more efficient for two of the three <i>C. species</i> when compared to free oil.</li> <li>EO-NE protect the components through nanoencapsulation and increase of the contact area due to the reduced size of the nanoemulsion may favor antibiofilm activity.</li> </ul>
[56]	<i>Cyclopia intermedia</i>	South Africa	Purchased	AQ ME fermented AQ	150 mg/mL 7.5 mg/mL 150 mg/mL	<i>C. albicans</i> . ATCC 10231;	ME show most effective against <i>C. albicans</i>
[57]	<i>Erythrina stricta Roxb.</i>	India	Stem bark	DCM EtOAc n-Hexane	7.8 mg/mL 125 mg/mL 125 mg/mL	<i>C. albicans</i> .	Both show significant antifungal activity against <i>C. albicans</i> . Present flavonoids and phenolics
[58]	<i>Matricaria recutita</i>	Egypt	-	Pharmacopeia (PhEur) grade essential oil	160 to 320 µg/mL.	<i>C. albicans</i> ATCC 90028	<ul style="list-style-type: none"> <li>Combination of essential oil with fluconazole and nystatin showed synergic inhibitory effects.</li> <li>Show the best when combining to tetracycline.</li> </ul>
[59]	<i>Piper hispidum</i>	Brazil	Leaves	Crude extract	62.5 mg/mL	<i>C. albicans</i> , <i>C. parapsilosis</i> <i>C. tropicalis</i> .	Show antifungal activity against <i>C. albicans</i> , by inhibition biofilm formation. Presents antimicrobial properties of chalcones
[60]	<i>Justicia glauca</i>	USA	Leaves	Green synthesis of gold nanoparticles (AuNPs) extract	12.5 ± 0.3 (µg/mL ± SD)	<i>C. albicans</i>	NPs greatly increased <i>J. glauca</i> against <i>C. albicans</i> by interference with growth-signaling pathway inside the cell via modulating tyrosine phosphorylation of growth essential peptides substrate
[61]	<i>Funtumia africana</i>	South Africa	Leaves	Isolated methyl ursolate HE CHL	Methyl ursolate: 63-µg/mL HE:40 µg/mL CH:80 µg/mL	<i>C. albicans</i> ATCC 10231	<ul style="list-style-type: none"> <li>CHL show strongest synergistic activities with methyl ursolate low toxicity which may support the use of this plant.</li> <li>Antimicrobial and anti-inflammatory activities of the crude extract provide some support for the traditional use of the plant.</li> </ul>
[62]	<i>Pappea capensis</i>	South Africa.	Leaves	HE DCM EtOAc BA extracts	0.39 - 0.78 mg/mL	<i>C. albicans</i>	Show antifungal activity
[63]	<i>Equisetum hyemale</i>	Japan	Stems	Crude extract DCM EtOAc	6.5 - 52.4 mg/mL	<i>C. albicans</i> <i>C. kefyr</i> <i>C. geochares</i> <i>C. krusei</i>	Show antifungal activity and negligible cytotoxicity. It contains epicatechin and β-carotene-linoleic acid

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, µg/mL)	Used Strains	Conclusion
[64]	<i>Croton limae</i>	Brazil	Leaves	Essential oil	>1024 (mg/mL)	<i>C. albicans</i> ATCC 40006 <i>C. krusei</i> ATCC 2538 <i>C. tropicalis</i> ATCC 40042	<ul style="list-style-type: none"> <li>Show antifungal activity against <i>C. albicans</i>, however antagonist effect was seen when combine with benzoyl-metronidazole.</li> <li>Cedrol, eucalyptol, a-pinene, b-pinene and linalool may be responsible for the antibacterial activity</li> </ul>
[65]	<i>Citrus sinensis</i>	USA	Peels	Essential Oils	1.68 µg /mL	<i>C. albicans</i> <i>C. tropicalis</i> <i>C. glabrata</i> <i>C. guilliermondii</i> , <i>C. lusitaniae</i> ,	<ul style="list-style-type: none"> <li>Moderate activity</li> <li>Can be used in oral hygiene products</li> <li>Less toxic alternative to amphotericin B.</li> </ul>
	<i>Citrus latifolia</i>				0.4 µg /mL		
[66]	<i>Haplophyllum tuberculatum</i> (Forssk.) A. Juss.	Tunisia	Leaves, stems	Essential oils	0.30 mg/mL	<i>C. albicans</i> ATCC 90028; <i>C. glabrata</i> ATCC 90030; <i>C. parapsilosis</i> ATCC 27853 <i>C. krusei</i> ATCC 6258.	<ul style="list-style-type: none"> <li>Effective and has potential to prevent cancer development.</li> <li>Significant correlation existed between the concentrations of the essential oils.</li> <li>Antifungal activity may be attributed to R-(+)-limonene, S(-)-limonene and octanol.</li> </ul>
[67]	<i>Camellia sinensis</i> (L.) O Kuntze	Brazil	Leaves	Green tea	16-33 µg/mL	<i>C. albicans</i> ATCC 14053 <i>C. albicans</i> ATCC 64548 <i>C. krusei</i> ATCC 6258	<ul style="list-style-type: none"> <li>Antifungal activity was highest in black tea&gt; green tea &gt;white tea</li> <li>Suggesting no direct relationship with the concentration of total phenols.</li> </ul>
				White tea	16-135 µg/mL		
				Red tea	>270 µg/mL		
				Black tea	16-33 µg/mL		
[68]	<i>Leucaena leucocephala</i>	Malaysia	Leaves	ME	3.15 - 25.0 mg/mL	<i>C. albicans</i> <i>C. tropicalis</i>	<ul style="list-style-type: none"> <li>Significant antifungal activity through inhibition of cell proliferation and induced apoptosis in MCF-7.</li> <li>Contained condensed tannins and phenols.</li> </ul>
[69]	<i>Trametes hirsuta</i> <i>Trametes gibbosa</i> <i>Trametes versicolor</i>	Serbia	Dried mycelia and fruiting bodies	EE	32.0 mg/mL	<i>C. albicans</i> BEOFB 811m <i>C. krusei</i> BEOFB 821m <i>C. parapsilosis</i> BEOFB 831m	Showed low antifungal potential in comparison with ketoconazole.
[70]	<i>Artemisia herba-alba</i>	Jordan	Aerials	Essential oils	1.25 mg/mL	<i>C. albicans</i> ATCC 10231 <i>C. parapsilosis</i> ATCC 90018 <i>C. tropicalis</i> ATCC 13803	Show antifungal and anti-inflammatory activities and without detrimental effects. Revealed an important inhibitory effect on germ tube formation
[71]	<i>Avicennia marina</i>	U.A.E.	-	EE	-	<i>C. albicans</i> SC5314	<i>L. inermis</i> and <i>P. oleracea</i> showed significant anti-C. activity and against biofilm formation Lower cytotoxicity and higher selectivity indices, both plant extracts represent promising area of future research.
	<i>Fagonia indica</i>				-		
	<i>Lawsania inermis</i>				10 µg/mL		
	<i>Portulaca oleracea</i>				10 µg/mL		
	<i>Salvadora persica</i>				25 µg/mL		
	<i>Asphodelus tenuifolius</i>				-		
	<i>Ziziphus spina-Christi</i>				-		

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, mg/mL)	Used Strains	Conclusion
[72]	<i>Aster yomena</i>	Korea	Aerial	ME and Isolated compound apigenin (AP)	ME: - AP: 2.5 µg/mL	<i>C. albicans</i> ATCC 90028	ME show no antifungal effect. Only isolated apigenin has the antifungal activity.
[73]	<i>Artemisia vulgaris</i>	Brazil	Leaves	Essential oils	100 µg /mL	<i>C. albicans</i> ATCC14057 <i>C. glabrata</i> ATCC2301 <i>C. krusei</i> ATCC6258 <i>C. parapsilosis</i> ATCC22018	All three are able to inhibit the growth of the <i>C.</i> genus yeasts. Differences in the contents of the chemical components in the essential oils significantly influence antifungal activity action.
	<i>Biden pilosa</i>				64 µg /mL		
	<i>Sphagneticola trilobata</i>				100 µg /mL		
[74]	<i>Muntingia calabura L.</i>	Philippines	Stem and dried leaf	EE	Leaf: 0.625 mg/mL Stem: 2.5 mg/mL	<i>C. albicans</i>	Show antifungal activity. Presence of sterols, flavonoids, alkaloids, saponins, glycosides and tannins
[75]	<i>Ixora megalophylla</i>	Thailand	Leaves stems	Petroleum ether (Pet), EtOAc EtOH	Leaf Pet:1250 mg/mL EtOAc:78 mg/mL EtOH:156 mg/mL	<i>C. albicans</i>	EtOAc extract from the leaves and the EtOAc and EE from the stems possessed antifungal activities.
					Stems Pet:1250 mg/mL EtOAc:78 mg/mL EtOH:78 mg/mL		
[76]	<i>Stegesbeckia orientalis</i>	China	-	EE, petroleum ether fraction (PE-SO), EtOAc, BA and water fraction (WE-SO).	EE:2.50 µg/mL PE-SO:4.0 µg/mL EtOAc :1.25 µg/mL BA:2.50 µg/mL WE-SO:2.50 µg/mL	<i>C. albicans</i> ATCC 1023	EE showed the strongest antimicrobial, antioxidant and cytotoxic activities.
[77]	<i>Berberis lycium Royle</i>	India	Roots	Berberine (BE), ME HE AQ	BE:41.6 ± 18.04 mg/mL ME: 187.5 ± 62.5 mg/mL HE: NA AQ: 8 mg/mL	<i>C. albicans</i> SKUAST- TAM-1	Pure berberine found more effective than crude extract, followed by methanolic and aqueous extracts.
[78]	<i>Calamus leptospadix Griff.</i>	India	Shoots	Saponin	80 mg/mL	<i>C. albicans</i> MTCC 3007	Significant amount of saponin possesses antimicrobial properties.
[79]	<i>Sapindus saponaria L.</i>	India	Trees	Hydro alcoholic extract	390-1560 µg/mL	<i>C. albicans</i> <i>C. glabrata</i> <i>C. tropicalis</i> <i>C. parapsilosis</i>	Show antifungal activity by acting on cell membrane causing cell lysis within 60min.
[80]	<i>Ziziphus nummularia</i>	India	Leaves	Zinc oxide nanoparticles (ZnO NPs) and leaf extract	>10mg/mL NP: 1.25mg/mL	<i>C. albicans</i> ATCC2091 <i>C. glabrata</i> NCIM3448	ZnO NPs exhibited very good antifungal activity, even better than standard antibiotic Amphotericin B attributed to the small size of synthesized ZnO NPs.
[81]	<i>Juniperus communis</i>	Portugal	Mature berries	Essential oils	0.039-0.16 %	<i>C. albicans</i> ESAB.	<ul style="list-style-type: none"> <li>Against all the tested organisms, support the use of traditional medication usage of this species.</li> <li>Inhibited by morphological changes in the cell membrane. Also, antimicrobial activity may due to monoterpenes, such as terpinen-4-ol and 1,8-cineol.</li> </ul>

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, mg/mL)	Used Strains	Conclusion
[82]	<i>Artemisia judaica</i> L.	Jordan	Aerial	Essential oil	1.25 µg/mL	<i>C. albicans</i> ATCC 10231 <i>C. parapsilosis</i> ATCC 90018 <i>C. tropicalis</i> ATCC 13803	<ul style="list-style-type: none"> <li>Significantly inhibited germ tube formation and disrupted preformed biofilms of <i>C. albicans</i>.</li> <li>It contains piperitone, camphor and ethyl cinnamate</li> </ul>
[83]	<i>Ipomoea procumbens</i>	Brazil	Leaves, stem and roots	Hydro-methanol extracts	>250 µg/ mL	<i>C. krusei</i> <i>C. parapsilosis</i> <i>C. albicans</i>	
[84]		Brazil	Leaves	Essential oil	4,096 µg/mL.	<i>C. albicans</i> <i>C. krusei</i> <i>C. tropicalis</i>	<ul style="list-style-type: none"> <li>The oil caused the inhibition of <i>C. albicans</i> and <i>C. tropicalis</i> by disrupting morphological transition.</li> <li>Related to selena-1,3,7(11)-trien-8-one and selina-1,3,7(11)-trien-8-one epoxide,</li> </ul>
[85]	<i>Cinnamomum verum</i>	Iran	Leaves and bark	Essential oils	125 to 175 mg/mL	<i>C. albicans</i> <i>C. Tropicalis</i> <i>C. Krusei</i> <i>C. Glabrata</i> <i>C. Parapsilosis</i> <i>C. Famata.</i>	Could be applied as supplementary agents along with conventional antifungal drugs.
	<i>Caryophyllium aromaticus</i>				700 to 1000 mg/mL		
	<i>Artemisia dracuncululus</i>				1000 to 2000 mg/mL		
	<i>Origanum vulgare</i>				173 to 350 mg/mL		
	<i>Cymbopogon citratus</i>				125 to 175 mg/mL		
[86]	<i>Baccharis trinervis</i> (Lam.)	Brazil	Aerial	Essential oil	X	<i>C. albicans</i> , <i>C. Parapsilosis</i> <i>C. Tropicali</i>	X
[87]	<i>Sedum sediforme</i>	Turkey	-	Petroleum ether (PE), AC ME	PE:8 ± 0.4 µg/mL AC:1 ± 0.2 µg/mL ME:1±0.3 µg/mL	<i>C. albicans</i>	<ul style="list-style-type: none"> <li>ME most active one.</li> <li>Contains 4 phenolic acids (protocatechuic acid, p-coumaric acid, caffeic acid, and chlorogenic acid and flavonoids(quercetin)</li> </ul>
[88]	<i>Mentha piperita</i>	Saudi Arabia	Aerial	Essential oil	1.50 ± 0.16 mg/mL	<i>C. albicans</i> ATCC 26790	<ul style="list-style-type: none"> <li>Show significant antifungal activity and potential to perform better than amphotericin B.</li> <li>Presence of high menthol and menthone components.</li> </ul>
[89]	<i>Curcuma aeruginosa</i> Roxb	Thailand	-	Essential oils	250 mg/mL	<i>C. albicans</i> ATCC 90028	<ul style="list-style-type: none"> <li>Major components are oxygenated monoterpenes, 1,8- cineole and camphor</li> </ul>
	<i>Curcuma glans</i> K. Larsen						
	<i>Curcuma cf. xanthorrhiza</i> Roxb						
[90]	<i>Thymus vulgaris</i>	Iran	-	ME	68 µg/mL	<i>C. albicans</i> ATCC10231	<ul style="list-style-type: none"> <li><i>C. Zeylanicum</i> show better antifungal activity compared to other.</li> <li>Antifungal activity may due to eugenol, cinnamic aldehyde, saponin, alkaloid and flavonoid.</li> </ul>
	<i>Caryophyllim aromaticus</i>				48 µg/mL		
	<i>Echinophora platyloba</i>				27 µg/mL		
	<i>Allium cepa</i>				75 µg/mL		
	<i>Cinnamomum zeylanicum</i>				18 µg/mL		

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, mg/mL)	Used Strains	Conclusion
[91]	<i>Cuminum cyminum</i> <i>Salvadora persica</i>	Iran	Seeds	Alcoholic extract	578 mg/L 4.9 mg/mL	<i>C. albicans</i> ATCC 14053 <i>C. dubliniensis</i> ATCC CD60, <i>C. glabrata</i> ATCC 90030 <i>C. krusei</i> ATCC 6258 <i>C. parapsilosis</i> ATCC 22019.	<ul style="list-style-type: none"> <li>Both show strong to moderate activity.</li> <li><i>C. cyminum</i> characterized by high amounts of <math>\alpha</math>-pinene, limonene and 1,8-cineole.</li> </ul>
[92]	<i>Paeonia lactiflora</i>	USA	Roots	EE	49 mg/mL 196 mg/mL	<i>C. albicans</i> SC5314 <i>C. albicans</i> ATCC 18804	Show antifungal activity associated with cell membrane integrity and permeability on (1.3)- $\beta$ -D-glucan synthase.
[93]	<i>Isodon flavidus</i> (Hand. -Mazz.)	China	Twigs and leaves	Crude extract	62.5 mg/mL	<i>C. albicans</i> .	Show antifungal activity <i>Flavin A</i> and <i>lophanic acid</i> can breakdown the formed biofilm of <i>C. albicans</i> .
[94]	<i>Rubus idaeus</i>	France	Ripe and unripe fruits	n-hexane, EtOAc BA	> 1000 $\mu$ g/mL	<i>C. albicans</i> <i>C. glabrata</i> <i>C. parapsilosis</i> .	<ul style="list-style-type: none"> <li>HE and EtOAc have significant anti-adhesion activity against <i>C. albicans</i></li> <li>Contains high condensed tannins.</li> </ul>
[95]	<i>Pogostemon heyneanus</i> <i>Cinnamomum tamala</i> <i>Camphor</i>	India	Leaves	Patchouli essential oil	0.6-1 mg/mL 0.6 mg/mL 1 mg/mL	<i>C. albicans</i> ATCC-90028 <i>C. Glabrata</i> MTCC 6507 <i>C. Tropicalis</i> MTCC 310	Inhibited the key virulent property of <i>C. Albicans</i> , the transition from yeast cells towards hyphal formation of <i>C. Albicans</i> .
[96]	<i>Pluchea dioscoridis</i>	USA	Leaf	EE	30 mg/mL	<i>C. albicans</i> strains	Exhibited high antifungal activity which cause changes in phospholipase, hemolysin, and secreted aspartyl proteinase gene expression could completely collapse the yeast cell and inhibit the growth.
[97]	<i>Equisetum telmateia</i>	Iran	Aerial	Superficial fluid extraction (SFE), cold maceration (CM) and Fractionation extracts (F)	SFE:32 mg/mL MC&F: >128 mg/mL	<i>C. albicans</i>	<ul style="list-style-type: none"> <li>SFE method show more appropriate for extraction against <i>C. albicans</i>.</li> <li>Antimicrobial attributed to the phenolic substances identified such as catechin, kaempferol derivatives and p-OH-benzoic acid.</li> </ul>
[98]	<i>Pogostemon cablin</i>	India	Purchased	Patchouli essential oil (PC, PH and PP)	PC: NA PH:25 mg/mL PP:50 mg/mL	<i>C. albicans</i>	PH exhibited better antifungal activities than the other two.
[99]	<i>Succisa pratensis</i>	Poland	Leaves or flowers	ME	0.11 mg/mL	<i>C. albicans</i> <i>T. mentagrophytes</i>	<ul style="list-style-type: none"> <li>Show antifungal activity.</li> <li>The compounds 10-(acetylmethyl)-(+), 3-carene, methyl linolenate, hexadecanoic acid, pentacosane, hexacosane, heptacosane and thymol having strong antimicrobial activity.</li> </ul>
[100]	<i>Tritomaria quinqueidentata</i> (Huds.)	China	-	Crude extract	>128 mg/mL	<i>C. albicans</i> wild strain SC5314 and four mutant strains DSY448, DSY653, DSY465, DSY654	Show antifungal activity

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, mg/mL)	Used Strains	Conclusion
[101]	<i>Citrullus colocynthis</i>	Iran	Fruits	Hydroalcoholic extracts	1.56 -12.5 mg/mL	<i>C. albicans</i>	Show antifungal activity
[102]	<i>Salvia rhytidia Benth(Mint)</i>	Iran.	-	ME.	3.125 to > 100 mg/mL	<i>C. albicans</i> <i>C. glabrata</i> <i>C. tropicalis</i> <i>C. krusei</i> <i>C. parapsilosis</i> <i>C. Lusitania</i> <i>C. guilliermondii</i>	Show antifungal activity
[103]	<i>Allium hushidari, Allium sativum</i>	Iran	-	Essential oil	0.25- 2 mg/mL	<i>C. albicans</i>	<ul style="list-style-type: none"> <li>Show antifungal activity.</li> <li>Antimicrobial activity is related to alkaloids, tannins, saponins, flavone and glycosides.</li> </ul>
[104]	<i>Laserpitium spp.</i>	Macedonia	Roots and rhizomes	Extract	1.25 mg /mL	<i>C. krusei</i>	<ul style="list-style-type: none"> <li>Inhibition of biofilm.</li> <li>Major components are isomontanolide, laserpitine and montanolide. All showed a more pronounced effect than fluconazole.</li> </ul>
[105]	<i>Anthemis nobilis, Foeniculum vulgare, Simmondsia chinensis, Nigella sativa, Trigonella foenumgraecum, Gadus morhua, Mentha piperita, Syzygium aromatic, Zingiber officinale</i>	Egypt	Purchased	Essential oils	Fennel oil :0.78% Others: NA	<i>C. albicans</i> <i>ATCC 10231</i> <i>C. Glabrata</i> <i>C. tropicalis</i>	<ul style="list-style-type: none"> <li>Fennel essential oil had significantly higher antifungal activities compared with other tested.</li> <li>Fennel essential oil alone or in combination with fluconazole could provide a promising approach in management of vulvovaginal candidiasis.</li> </ul>
[106]	<i>Cocos nucifera</i>	Brazil	Purchased	NaP	6.25 µg/mL	<i>C. albicans</i> <i>C. glabrata,</i>	<ul style="list-style-type: none"> <li>Exhibited high antifungal activity against pathogenic <i>Candida</i> spp.</li> <li>The nano-capsules formulations prolonged storage, and increased photostability of clotrimazole and prolonged drug release.</li> </ul>
[107]	<i>Lycium barbarum</i>	Romania	Leaves	Phenolic oil	0.031-0.062 mg/mL	<i>C. albicans</i> <i>ATCC 10231,</i> <i>C. parapsilosis</i> <i>ATCC 22019</i>	<ul style="list-style-type: none"> <li>Show antifungal activity.</li> <li>The leaves contain higher amounts of chlorogenic acids and flavonoid glycosides.</li> </ul>
[108]	<i>Garcinia xanthochymus</i>	Brazil	Fruits	Xanthochymol and garcinol, isoprenylated benzophenones	1 to 3 µg/mL	<i>C. albicans</i>	<ul style="list-style-type: none"> <li>Show antifungal activity and can also potentiate the activity of fluconazole.</li> <li>inhibited development of hyphae and subsequent biofilm maturation, inducing cell death</li> </ul>
[109]	<i>Spondias tuberosa</i>	Brazil	Leaves	HE	2.0 mg/mL	<i>C. albicans</i> URM 5901, from unguai scales	<ul style="list-style-type: none"> <li>Show antifungal activity by disrupting cell membrane.</li> <li>It contains flavonoids, hydrolysable tannins, saponins, terpenes and unsaturated fatty acids</li> </ul>

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, mg/mL)	Used Strains	Conclusion
[110]	<i>Holothuria scabra</i> , <i>Holothuria parva</i> , <i>Holothuria leucospilota</i>	Iran	-	Crude extract	NA	<i>C. albicans</i> ATCC 10231	X
[111]	<i>Glycyrrhiza glabra</i>	Spain	Rhizomes and roots	Phenolic extract	>1.5 mg/mL	<i>C. glabrata</i> , <i>C. parapsilosis</i> <i>C. albicans</i> .	X
[112]	<i>Myracrodruon urundeuva</i>	Brazil	Bark	EE	4-512 µg/mL (topical)	<i>C. albicans</i> <i>C. krusei</i> <i>C. tropicalis</i>	Show antifungal activity which contains flavonoids and tannins.
[113]	<i>Ficus elastica</i> Roxb. ex Hornem.	Cameroon	Aerial roots	ME	4.9 µg/mL;	<i>C. albicans</i>	ME show antifungal activity. The most active antimicrobial components are elastiquinone and fucusosid
[114]	<i>Daucus virgatus</i> (Poir.) Maire	Tunisia	Aerial	EtOAc ME	625 µg /mL	<i>C. albicans</i>	Exhibited moderate activity
[115]	<i>Pistacia vera</i> L., Bronte	Italy	Hulls	Essential oil	2.5-5 mg/mL	<i>C. albicans</i> ATCC 64550 4 clinical strains of <i>C. albicans</i> ,	Show antifungal activity.
[116]	<i>Anisophyllea laurina</i> R. Br. ex	Guinea	Pulp seed	ME EE	500-1000 µg/mL	<i>C. albicans</i>	Both ethanol and methanol are very effective to extract phenolics and show antifungal activity.
[117]	<i>Thymus vulgaris</i> , <i>Citrus limonum</i> , <i>Pelargonium graveolens</i> , <i>Cinnamomum cassia</i> , <i>Ocimum basilicum</i> , <i>Eugenia caryophyllus</i>	Poland	Purchased	Essential oils	Cinnamon oil: 0.002–0.125% (v/v) Others: 0.005% or less to 2.5% (v/v).	<i>C. albicans</i> and 76 isolates of <i>C. glabrata</i>	<ul style="list-style-type: none"> <li>Cinnamon oil is the most active against <i>C. albicans</i>.</li> <li>All of the tested oils demonstrated the ability to inhibit the transition of yeast to mycelium form. Thyme, lemon, and clove oils affected cell membranes by influencing potassium ion efflux, which was not seen in the lemon oil.</li> <li>No synergistic interactions between antifungal drugs; possible synergism was between amphotericin B and geranium oil.</li> </ul>
[118]	<i>Lippia sidoides</i> Cham	Brazil	Purchased	Essential oil	156 and 312 µg/mL	<i>C. albicans</i> ATCC 64548	<ul style="list-style-type: none"> <li>Show antifungal activity against <i>C. albicans</i></li> </ul>
[119]	<i>Mentha piperita</i>	Brazil	Purchased	Essential oil	1.875 µg/mL	<i>C. albicans</i> INCQS 40277 <i>C. tropicalis</i> ATCC 28707	Show antifungal activity and inactivate potentially spoilage yeasts in fruit juices through disturbance of different physiological functions in yeast cells.
[120]	<i>Hippophae rhamnoides</i> L	Poland	Twigs and leaves	Extract	250 mg/mL (twig), 31.5 mg/mL (leaf)	<i>C. albicans</i> , ATCC 10231 <i>fluconazole-sensitive</i> and <i>clinical</i> , <i>C. glabrata</i> G1	Significant antifungal activity by inhibited morphogenesis such as germ tube and hyphae formation.
[121]	<i>Paeonia lactiflora</i>	Korea	Root	EE	196 µg/mL	<i>C. albicans</i> , ATCC 188040 <i>C. albicans</i> KCCM 50235	EE show good inhibitory effects against biofilm formation by impeding cell adhesion and obstructing the morphological transition of hyphae. Also inhibited the cell wall synthesis and damages cell membrane functions which lead to cell swelling and lysis.

(Table 3) Contd....

Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, mg/mL)	Used Strains	Conclusion
[122]	<i>Psidium guajava</i>	Brazil	Leaves	AQ and Hydroethanolic extract	>8192 mg/mL	<i>C. albicans</i> <i>C. tropicalis</i>	X
[123]	<i>Kedrostis africana</i>	South Africa	Dried tubers	AcE AQ EE	Ac:0.312 mg/mL Aq: >5 mg/mL Eth:0.325 mg/mL	<i>C. albicans</i> ATCC 10231	<ul style="list-style-type: none"> <li>Show antifungal activity.</li> <li>The flavonoid, proanthocyanidin and total phenolic concentrations were higher in AcE compared to the aqueous and EE.</li> </ul>
[124]	<i>Eucalyptus microcorys</i>	Australia	Leaves	AQ	1250 µg/mL.	<i>C. albicans</i> ATCC 10231	<ul style="list-style-type: none"> <li>Show antifungal activity and was found to be a good source of TPC, TFC, proanthocyanidins and saponins.</li> </ul>
[125]	<i>Psiadia punctulata</i> (DC.) Vatke	Saudi Arabia	Leaves	Extract	50 µg/mL	<i>C. albicans.</i>	<ul style="list-style-type: none"> <li>Show antifungal activity against <i>C. albicans</i>.</li> <li>Isolated 3',4',5,7-tetramethoxyflavone, displayed the ability to reduce biofilm formation of <i>C. albicans</i> by 90%</li> </ul>
[126]	<i>Camellia sinensis</i> (L.) O Kuntze	Korea	Seed	Green tea seed extract	938 µg/mL	<i>C. albicans</i> ATCC 10231	<ul style="list-style-type: none"> <li>Active compounds: theasaponin E1, assamsaponin A and assamsaponin B</li> <li>GTS extract can be used as a safe and strong natural anti-yeast.</li> </ul>
[127]	<i>Psidium guajava</i> , <i>Psidium brownianum</i> Mart. Ex DC	India	Leaves	Hydroethanolic extracts	8,192 µg/mL,	<i>C. albicans</i> <i>C. tropicalis</i> strains	<ul style="list-style-type: none"> <li>Show antifungal activity against <i>C. albicans</i> and are effective on potentiating the effect of fluconazole.</li> <li>Presence of phenols, flavonoids and tannins.</li> </ul>
[128]	<i>Murraya koenigii</i> (L.) Spreng	India	Leaves	Hydro-distillate essential oil	12.5-100 µg/mL	<i>C. albicans</i> strain MTCC 3017	Show antifungal activity against <i>C. albicans</i> inhibited by the compound mk309
[129]	<i>Melaleuca alternifolia</i>	-	-	Essential oils	0.25-2% v/v	<i>C. albicans</i> ATCC 10231, <i>C. glabrata</i> , <i>C. tropicalis</i> , <i>C. parapsilosis</i> <i>C. krusei</i> , <i>C. guilliermondii</i> , <i>C. lusitaniae</i> , <i>C. dubliniensis</i> ,	<ul style="list-style-type: none"> <li>All strains show antifungal activity</li> <li>Peppermint oil demonstrated the lowest antifungal activity.</li> </ul>
	<i>Mentha piperita</i>				0.03-0.25% v/v		
	<i>Thymus vulgaris</i>				0.25-2% v/v		
	<i>Syzygium aromaticum</i>				0.06-0.25 % v/v		
[130]	<i>Combretum erythrophyllum</i>	South Africa	Leaves	AQ AcE DCM HE	1.25 mg/mL	<i>C. albicans</i>	<ul style="list-style-type: none"> <li>Antifungal activity followed by AQ &gt; AcE &gt; DCM &gt; HE</li> <li>Provide some indication for the traditional use of the plant.</li> </ul>
[131]	<i>Strychnos spinosa</i> Lam.	Nigeria	Leaves	AcE ME DCM	1.25 or >1.25 mg/mL	<i>C. albicans</i> ATCC 10231	Show antifungal activity and support the traditional use of this plant as treatment of infectious.
[132]	<i>Aloe trigonantha</i> L.C. Leach	Ethiopia	leaf latex	Aloesin, 8-O-Methyl-7-hydroxyaloin	400 µg/mL.	<i>C. albicans</i> ATCC 10231	Show weak antifungal activity

(Table 3) Contd....




Refs.	Plant/Organism	Country of Origin	Plant Part(s)	Formulation	MIC (mg/mL, µg/mL)	Used Strains	Conclusion
[133]	<i>Nigella sativa</i>	India	Leaves	Hydro-steam distilled essential oils	15.62 µg/mL	<i>C. albicans</i> MTCC-183, <i>C. tropicalis</i> MTCC-184, <i>C. glabrata</i> MTCC-3019	Ajwain and Black Cumin leaf oils showed better antifungal activity by inhibition of cell membrane synthesis, specifically by extracting the sterols from the membrane or inhibiting steroid synthesis.
	<i>Murraya koenigii</i>				250 µg/mL		
	<i>Trachiyspirum ammi</i>						
	<i>Piper betel</i>						
[134]	<i>Carpolobia lutea</i>	Nigeria	Leaves	EE ME AQ HE	25 mg/mL	<i>C. albicans</i>	<ul style="list-style-type: none"> <li>• EE show significant antifungal effect.</li> <li>• Both AQ and HE show no inhibitory effect.</li> </ul>
[135]	<i>Zuccagnia punctata</i> Cav.	Argentina	Aerial parts	DCM	27.33-31µg/mL	<i>C. albicans</i> <i>C. glabrata</i>	<ul style="list-style-type: none"> <li>• Show antifungal activity. <i>ZpE-LnE</i> have synergistic effect, support the proper joint use of both antifungal herbs in traditional medicine.</li> </ul>
	<i>Larrea nitida</i> Cav.						
[136]	<i>Tetraglochin cristatum</i> (Britton) Roth	Argentina	Leaves and aerial	Hydroalcoholic dry extract	12.5 and 25 µg/mL	<i>C. albicans</i> 144783, 134333, 2089; <i>C. glabrata</i> 031646, 042030, 031982; <i>C. tropicalis</i> 1841;	<ul style="list-style-type: none"> <li>• Show antifungal activity and give support to their traditional use for treating infections.</li> <li>• It contains hydrolysable and condensed tannins</li> </ul>
[137]	<i>Satureja Khuzistanica</i>	Iran	Aerial	EE	299.4 mg/mL	<i>C. albicans</i> ATCC 10231 <i>C. albicans</i> ATCC 66506 a	Show synergistic effect with amphotericin B and ketoconazole, while this extract had no effect on clotrimazole activity.
[138]	<i>Alchemilla vulgaris</i> L.	Serbia	Root	ME	>20 µg/mL	<i>C. albicans</i> ATCC 10259,	X
[139]	<i>Ferula assafoetida</i>	Iran	oleo-gum-resin	Essential oil	0.19 (0.12-0.25) µg/mL	<i>C. albicans</i> CBS 5982, 1905 and 1949	Show remarkable antifungal activities
[140]	<i>Thymus vulgaris</i>	Brazil	Leaves	Extracts	50 mg/mL	<i>C. albicans</i> ATCC 18804, <i>S. aureus</i> ATCC 6538	Show antifungal activity by acting on the biofilm formation. It contains thymol, carvacrol, linalool, geraniol, citral, tannins, organic acids, flavonoids, minerals.
[141]	<i>Eugenia leitonii</i> , <i>Eugenia brasiliensis</i> , <i>Eugenia myrcianthes</i> <i>Eugenia involucrata</i>	Brazil	Leaves pulp seeds barks	Dry extracts	Barks: 15.62 ->2000 µg/mL <i>E. leitonii</i> (seed): 15.62 µg/ mL <i>E. brasiliensis</i> (leaf): 31.25 µg/ mL <i>E. brasiliensis</i> (seed): 5.62 µg/ mL	<i>C. albicans</i> ATCC 90028	The seeds of <i>E. leitonii</i> and the seeds and leaves of <i>E. brasiliensis</i> were found to have strong antifungal activity against <i>C. albicans</i> by acting on mature biofilms. However, Bark show no antifungal effect. Phenolic compounds epicatechin and gallic acid were the major constituents in the extracts.

Abbreviations: \* -: Not specified/Not available, \*X: No antifungal effect.; \*ME: Methanolic extract, \* AC: Acetone extract, \*AQ: Aqueous extract, \*AQE: Aqueous ethanolic extract; \*EtOAc: Ethyl acetate extract, \* EE: Ethanol extract, \*Dichloromethane extract: DCM, \*HE: Hexane extract, \*CHL: Chloroform extract, \*BA: butanol extract, \*NaP: Nanoparticles/Nano formulation.

The most common source types investigated were the aerial parts of the plants. Methanolic and ethanolic extractions exhibited higher antifungal efficacy, amongst other extracts. A total of 30 articles investigated the mechanisms of the herbal extracts against *C. albicans*, which they exert antifungal effects through inhibiting biofilm formation, hyphal transformation, germ tube inhibition; alteration of

membrane potential and permeability; disrupting transcription, cell division and inhibition of virulence factors.

Overall, the presented evidence shows that natural products may be employed effectively as an alternative therapy against *C. albicans*. Among the tested plants, common plants with a long history and well-known beneficial effects such as




**Fig. (1).** Flowchart of search strategy and study selection procedure. (A higher resolution/colour version of this figure is available in the electronic copy of the article).

tea (*C. sinensis*), tea tree oil (*M. alternifolia*), cinnamon (*C. verum*), cumin (*C. cyminum*), henna (*L. inermis*), mint (*M. piperita*), and thyme (*T. vulgaris*), whose antifungal activity was confirmed in several studies of this review, support the traditional use of these plants. Furthermore, several novel natural herbs have been discovered as potential adjunctive treatments against *C. albicans*.

### 3.2. Herbal Interventions *In Vivo*

When investigating the *in vivo* effects of herbal extracts on *C. albicans*, 9 articles that matched the search criteria with a total of 11 plants were examined (Table 2). Most studies used rats as the host organism by infecting them with *C. albicans* in which, only *Vicia faba* and *Morinda tomentosa* show no antifungal activity against *C. albicans* [12, 14]. The



**Fig. (2).** Distribution of the geographical locations of the origins of plants cited in this review, per source country. (A higher resolution/colour version of this figure is available in the electronic copy of the article).

most active plants were *C. sinensis* with MIC values of 40  $\mu\text{g/mL}$  [13]. Topical applications of 4% *L. innermis* show similar or better effect than clotrimazole and *Mitracarpus frigidus* presents antifungal activities greater than fluconazole against candidiasis [11, 16]. *M. alternifolia* inhibited biofilm formation [15] *Syzygium cumini* and *Jatropha curcas*, which contained a high amount of phenols and flavonoids are responsible for the inhibitory effect against *C. albicans* [17, 19]. Nano-formulations of the seeds of *S. cumini* exhibited an improved antioxidant activity of the plant extract as compared to other formulations [17]. When comparing all the *in vivo* herbal interventions, *L. innermis* and *M. alternifolia* seem to have the most significant antifungal activity with the MIC values as low as 5-10 mg/mL *in vivo* and CFU value 5.33 Log<sub>10</sub> [10, 15]. Nano formulations of all seven herbal extracts that were included in this review (*P. graveolens*, *Eucalyptus globulus*, *Justicia glauca*, *Ziziphus nummularia*, *Pogostemon cablin* and *Cocos nucifera*) demonstrated increasing plants properties and better antifungal effects than traditional extraction methods (Tables 2 and 3).

### 3.3. Herbal Interventions *In Vitro*


Table 3 summarizes the herbal interventions *in vitro*, where 122 studies were included explaining the effects of 175 herbal interventions [20-141]. Most of the herbal extracts examined demonstrate activities against *C. albicans* in which, the most active were *C. sinensis*, *Citrus sinensis*, *Citrus latifolia*, *C. nucifera*, *Ficus elastica* Roxb. Ex Hornem, *M. piperita*, *Garcinia xanthochymus*, *M. alternifolia*, *P. graveolens*, *Siegesbeckia orientalis*, *Sedum sediforme* and *L. innermis* with MIC values ranging from 0.0945  $\mu\text{g/mL}$  to 10  $\mu\text{g/mL}$ . A total of 25 natural extracts were ineffective against *C. albicans*. Twelve articles assessed the synergistic effect of

herbal extracts with antifungal drugs in which, *M. alternifolia*, *Buchenavia tetraphylla*, *Foeniculum vulgare*, *G. xanthochymus*, *P. guajava*, *Psidium brownianum* Mart. ex DC and *Satureja khuzistanica* showed a synergistic effect with fluconazole. *Matricaria recutita* showed an additive effect with nystatin and fluconazole. *P. graveolens* showed synergism with amphotericin B. *S. khuzistanica* potentiated the effect of amphotericin B and ketoconazole. However, five tested extracts, which are *T. vulgaris*, *Citrus limonum*, *Cinnamomum cassia*, *Ocimum basilicum* and *Eugenia caryophyllu* showed little or no enhancement. The antagonist effect of *Croton limae* with benzoyl metronidazole was observed.

Three herbal interventions demonstrated greater antifungal effect than common antifungal drugs in which, *Z. nummularia* presented better antifungal activities over Amphotericin B, due to the incorporations of synthesized zinc nanoparticles that help to enhance plant properties [80]. The presence of isomontanolide, lesertipine and montanolide in *Laserpitium* species exhibited a more pronounced effect than fluconazole by inhibiting the hyphae and subsequent biofilm maturation [104]. In a study of the aerial parts of *M. piperita*, the MIC value of the extracts (1.5  $\mu\text{g/mL}$ ) [88] was smaller than the Amphotericin B (MIC: 5  $\mu\text{g/mL}$ ), implying that it has the potential to perform better antifungal activities than the synthetic drugs.

### 3.4. Preparation of Herbal Extracts

Differences in parts of plants extracted and extraction methods on the same plant greatly influenced its antifungal activities. In the study of *Piper guineense*, the fruits and leaves were prepared in five different extracts. Among these, aqueous extraction shows no inhibitory effect, while methanolic



**Fig. (3).** Chemical structures of (a) Catechins, (b) Gallic acid, (c) Thymol, and (d) condensed Tannins, as described in the systematic review.

extracts present the most significant activity with MIC of 39  $\mu\text{g/mL}$  and the rest with MIC of 78  $\mu\text{g/mL}$  [24]. In another study investigating Leguminosae family species, the authors examined the leaves of 8 plants which were prepared in both methanolic and ethanolic extracts, showing that *Withania somnifera*, *Echinophora platybola* and *Zingiber officinale* demonstrated better effect in ethanolic extracts, while *Curcuma longa* and *Pogostemon parviflorus* present better effect in methanolic extracts [28]. Another study, using hexane and chloroform extracts of the leaves of *B. tetraphylla*, exhibited significantly greater inhibition as compared to ethanolic and methanolic extracts [48]. It appears that it is difficult to generalise and select an extraction method that preserves the greatest activity of the plant extract that is valid for all plants. It is equally difficult, given the range of plant types and parts used in the studies, to generalise about the specific part of a plant, which would give the highest concentration of the active substance. It can, therefore, be concluded that the extraction method and selection of the part of the plant to be used must be individualised on a plant-by-plant basis.

However, when comparing common extraction methods such as methanol and aqueous extracts with nano-formulated extracts, the use of nanotechnology greatly improves the plant's properties and the antifungal effects remain active for a longer period of time. The increase in antifungal activity may be attributed to its small size (200nm), which may activate the passive transport mechanism across the cell membrane. The study of *P. graveolens*, revealed that nano-formulations of geranium oil (MIC: 1.82  $\mu\text{g/mL}$ ) are twice as active as crude extracts (MIC: 3.64  $\mu\text{g/mL}$ ) and the former are sufficiently active to reduce the amount of biofilm on catheters [117]. The nanoparticles protect the components

through nanoencapsulation and increase the contact area due to the reduced size of the formulations, which improves anti-biofilm activity. In the study of *J. glauca* against *C. albicans*, the nanoparticles greatly inhibited bacterial growth by interfering with the growth-signaling pathway inside the cell via modulating tyrosine phosphorylation of growth essential peptides substrate [60]. In addition, in the study of *C. nucifera*, the nano-capsule formulations exhibit favourable properties after 60 days of storage and in prolonged drug release [106].

Several studies have examined the significant antifungal effects of these herbs more than once, which are *S. persica*, *C. sinensis*, *C. verum*, *E. uniflora*, *L. inermis*, *M. alternifolia*, *M. piperita*, *P. lactiflora*, *P. graveolens*, *S. cumini* and *T. vulgaris*. Two studies [71, 91] revealed that *S. persica* (MIC: 25  $\mu\text{g/mL}$  and 4.8mg/mL) had strong to moderate activity against different pathogenic *Candida* species. Both use the alcoholic extraction methods and the results are in accordance with each other.

Three studies demonstrated the activity of *C. sinensis* [13, 58, 118] in which, the leaf extracts (MICs ranging from 16-135  $\mu\text{g/mL}$ ) exhibited higher antifungal activities than the seeds (MIC: 938  $\mu\text{g/mL}$ ). Also, studies revealed that green and black leaves present better activities than white and red tea leaves, which may be due to different fermentation methods. What's more, a higher percentage of catechins are found in green tea leaves, which are well known for their antioxidant activity. Catechins are reducing agents or chelating metal ions, which are able to inhibit both DNA damage and lipid peroxidation, ultimately cause membrane integrity [142]. The final study showed that *C. sinensis* was effective

both *in vivo* and *in vitro* against *C. albicans* [13]. All three studies demonstrate a significant effect of tea tree against *C. albicans*.

When comparing two studies using cinnamon [37], the aerial parts of the plant with the MIC values of 31.25 to 62.5 µg/mL exhibited greater fungicidal effects than the leaves and bark (MIC: 127-175 µg/mL) [95]. Both studies revealed the importance of cinnamaldehyde and cinnamaldehyde dimethyl-acetate against microorganisms. In other studies, all five studies investigating *T. vulgaris* exhibited significant antifungal activities of this plant against *C. albicans* [50, 91, 118, 129, 140]. The results demonstrated the importance of thymol as an active agent inhibiting biofilm formation, promoting high cell viability, having anti-inflammatory effects and presenting no genotoxicity [140].

### 3.5. Active Compounds

Phenolic compounds have been studied extensively of their wide range of antioxidants and beneficial effects on the human body for decades. In this systematic review, numerous active compounds have been identified to be active against *C. albicans*. Compounds that stand out for their marked antifungal activity include phenols such as gallic acid, thymol, and flavonoids (especially catechin – Fig. 3a), polyphenols such as tannins, terpenoids and saponins.

Gallic acid is a trihydroxybenzoic acid with antioxidant, anti-inflammatory, and antimicrobial properties (Fig. 3b). In this review, four articles reveal the antifungal effectiveness of gallic acid [17, 31, 46, 141]. Particularly in the study of *Cochlospermum regium*, it has been demonstrated that the antifungal mechanism of gallic acid is either by binding to ergosterol on the cell membrane that leads to pore formation or by disrupting the enzymes responsible for the ergosterol synthesized, thereby causing membrane damage [32].

Thymol (2-isopropyl-5-methylphenol) isomeric with carvacrol (Fig. 3c) is the main monoterpene phenol isolated from plants belonging to the *Lamiaceae*, *Verbenaceae*, *Scrophulariaceae*, *Ranunculaceae*, and *Apiaceae* families. It has been used for treatment due to its antioxidant, anti-inflammatory, local anaesthetic, antinociceptive, antiseptic, antibacterial, and antifungal effects as well as for their beneficial effects on the cardiovascular system [143]. The studies of *T. vulgaris* and *Succisa pratensis* revealed that thymol is able to block ergosterol synthesis and ultimately caused pore formation in the membrane [49,99].

Tannins (Fig. 3d) are known for their potent antioxidant, cytotoxic and antimicrobial activities. The study of *Ricinus communis* suggests that the potential fungicidal activity occurs by the targeting of surface-exposed adhesins, cell wall polypeptides and membrane-bound enzymes of the fungal cell. The proposed mechanism involves complex formation between tannins and flavonoids with nucleophilic amino acids in proteins, leading to the inactivation of the proteins and loss of function. The methanolic extracts show greater antifungal effective over aqueous and ethanolic extracts, due to the higher preservation of the tannins, flavonoids and terpenoid compounds in the extracts. In this systematic review, tannins were found to be present and showed antifungal ac-

tivity in 11 herbal extracts [18, 31, 47, 75, 95, 104, 109, 113, 128, 136, 140].

Saponins are phytochemicals, which can be found most in peas, soybeans and herbs. In this review, multiple studies reveal the potential antifungal activities against *C. albicans* in the presence of saponins [40, 41, 75, 79, 80, 91, 104, 110, 125, 127]. However, in this review, the detailed mechanisms of action of saponins have not been well investigated. Previous research demonstrated that saponin is able to interfere with sterols, leading to inhibition of yeast-hyphal transition and biofilm formations [144].

Flavonoids are metabolites widely present in most vegetables, particularly green and red vegetables. Traditional and local communities used these plants due to their anti-inflammatory, antioxidant, anti-depressant and anti-infective effects. The mechanism of action has not been elucidated completely, even though it is believed to interfere with the cell wall and/or the ergosterol synthesis [145]. In this present review, several articles exhibited the importance of flavonoids in against microorganisms [17, 19, 21, 31, 40, 43, 47, 58, 75, 88, 91, 108, 110, 113, 124, 128, 140]. In the study of *S. cumini*, stating that plants containing high flavonoids were found to have strong inhibitory effects on the formation and metabolic activity of *C. albicans* biofilms or planktonic cells [40]. Catechin is a flavan-3-ol type natural phenol commonly found in oolong and green tea. Its anti-oxidant, anti-hypertensive, anti-inflammatory, anti-proliferative, anti-thrombogenic, and anti-hyperlipidemic activities have been clearly illustrated through various *in vitro* and *in vivo* studies. It was found that catechin can induce the generation of reactive oxygen species (ROS). ROS are implicated in the disruption of molecular mechanisms such as angiogenesis, extracellular matrix degradation and have been shown to lead to cell apoptosis [146]. The antifungal effect of catechins is demonstrated across several of the articles included in this review, all supporting the role played by catechin as an antifungal against *C. albicans* [47,68,98,141]. In addition to catechin, green tea seeds extract contain theasaponin E1, assamsaponin A and assamsaponin B, all of which were active against *C. albicans* and may have applications in food preservation against yeast contamination [127].

Terpenoids, sometimes called isoprenoids, can be found in the leguminous plant, turmeric and mustard seed. In this review, a number of investigations report that plant extracts like *Helichrysum* and *Juniperus communis* containing terpenoids exhibited antifungal activity against *C. albicans* [22, 31, 45, 82, 90, 110]. The proposed mechanism of action is that terpenoids have a fungistatic effect on *Candida* by modulating specific signaling pathways (TOR pathway or calcium signalling), rather than by creating nonspecific membrane lesions. The result of this is the alteration of gene transcription and stasis [147].

Other similar active compounds have been identified, such as g-terpinene and 1,8-cineole in *Lavandula binaludensis* and *C. cyminum* [37,91]; 5-O-methylatfolin; epilupeol acetate in *Ficus drupacea L.* [42];  $\alpha$ -pinene, aromadendrene, globulol, betulinic acid, oleanolic acid-3-acetate and ursolic acid-3-acetate present in *S. glomulifera*, *C. cyminum* and *S. persica* [49];  $\alpha$ -pinene, limonene and 1,8-cineole, oleanolic

acid and luteolin-7-*O*-glucoside in *Haplophyllum tuberculatum* (Forssk.) A. Juss. [66]; vepicatechin and  $\beta$ -carotene-linoleic acid in *Equisetum hyemale* [63]; and selenal,3,7(11)-trien-8-one in *E. uniflora* [63]. Further studies are required in order to characterize their antifungal activity against *C. albicans*, since their mechanisms of action are not yet well established.

## 2.6. Strengths

Only clinical studies, clinical trials and RCTs are included in this systematic review in order to reduce experimental bias. Most trials used placebos or standard antifungal agents in the control group. Overall, the majority of herbal interventions reviewed indicate the antifungal effect and the major bioactive compounds responsible for the antifungal activity against *C. albicans*.

## 2.7. Limitations

This systematic review highlights the lack of consensus and standardization of MIC values defining the strength of antifungal activity specifically for *C. albicans*. Each investigator and study determine its own scale regarding what is significant inhibition and what is not. For example, the studies of *Glycyrrhiza glabra* (MIC:1500  $\mu$ g/mL) and *Rhaphiodon echinus* (MIC:1024  $\mu$ g/mL) are reported inactive [11, 46], on the contrary, in the case of *M. tomentosa* with MIC value as low as >32  $\mu$ g/mL is considered ineffective by the investigator [14]. However, in general, most authors considered MIC values below 100  $\mu$ g/mL as significant; between 100-1000  $\mu$ g/mL as moderate; and above 1000  $\mu$ g/mL as inactive.

The majority of studies utilized methanolic and ethanolic extracts, whereas other extraction methods such as aqueous, hexane, ethyl-acetate, acetone and dichloromethane. Other formulations studies used essential oils. Some drawbacks of essential oils have been identified, such as chemical complexity, high volatility, susceptibility to degradation and oxidation, insolubility in aqueous systems and low bioavailability [148]. These characteristics hinder their direct use of products, although all studies established the extraction process and methods thoroughly and in detail. Several factors, like temperature, pH, particle size and solvent, may affect the outcomes.

Furthermore, the difference in concentrations, quantities, incubation time and treatment duration are not equivalent, which will greatly influence the outcomes and make it difficult to compare. Most of the *in vitro* studies are incubated mostly for 24 hours; however, in some studies, it may extend to 48 to 72 hours, allowing the formation of the biofilm. The incubation temperatures range from 30-37°C, and a variety of culture methods are used across the studies, for example, agar, microwell, and sabouraud dextrose agar plates. As for the treatment duration, *in vivo* studies using *C. sinensis* and *M. alternifolia*, the mice were treated for 5 days, whereas, in the study using *S. cumini*, the rats were treated for 21 days [13, 15]; and the treatment period was 2 weeks in the study of *Astragalus membranaceus* [18]. Several herbal and extraction solvent concentrations were used in which it can be

postulated that a longer period of treatment and higher concentrations could lead to an overestimation of positive outcomes.

The parts of the plants collected are also different such as bark, roots, leaves, flowers, hulls and seeds. A difference in concentrations used between *in vivo* and *in vitro* studies could lead to variation in response mechanisms towards the extracts. One drawback relates to this review is that majority of the studies were the very first study of examining the activity of the extracts or in the early phase of trials. Several studies did not illustrate the antifungal mechanisms and active components. Another concern is that the safety measure of the studies and their potential interactions with other drugs were not investigated. Further studies are needed to ensure the effectiveness, determine the mechanisms of action as well as efficacy, safety and intrinsic toxicity of the active compounds *in vivo*. During the data collection process, only English studies are included; therefore, language bias could be another restriction of this study.

## CONCLUSION

The results show that a wide range of plant extracts are able to inhibit *C. albicans* *in vitro*. The most active extracts were *M. alternifolia*, *Cit. sinensis*, *C. latifolia*, *C. nucifera*, *F. elastica* Roxb. Ex Hornem, *M. piperita*, *G. xanthochymus*, *P. graveolens*, *Sedum sediforme*, *L. inermis* and *S. orientalis*. The least active extracts were *R. echinus*, and *G. glabra*. The most active extract *in vivo* was *C. sinensis*.

The most common source types investigated were the aerial parts. Most plants with methanolic and ethanolic extraction exhibited high antifungal efficacy, amongst others. This could be due to the increased solubility of non-polar compounds in such solvents [151]. The most crucial components that have proved to have antifungal activities were the phenols such as gallic acid, thymol, and flavonoids (especially catechin), polyphenols such as tannins and terpenoids. The incorporation of nanotechnology shows promising results in the use of natural compounds against bacterial infections.

## LIST OF ABBREVIATIONS

DNA	=	Deoxyribonucleic acid
RNA	=	Ribonucleic acid
BCE	=	Before Common Era
CFU	=	Colony Forming Units
MIC	=	Minimum Inhibitory Concentration
TTO	=	Tea Tree Oil
PRISMA	=	Preferred Reporting Items for Systematic Reviews and Meta-Analyses

## CONSENT FOR PUBLICATION

Not applicable.

## FUNDING

None.



## CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

## ACKNOWLEDGEMENTS

Declared none.

## REFERENCES

- [1] Spampinato, C.; Leonardi, D. *Candida* infections, causes, targets, and resistance mechanisms: traditional and alternative antifungal agents. *BioMed Res. Int.*, **2013**, 2013204237 <http://dx.doi.org/10.1155/2013/204237> PMID: 23878798
- [2] Sardi, J.C.O.; Scorzoni, L.; Bernardi, T.; Fusco-Almeida, A.M.; Mendes Giannini, M.J.S. *Candida* species: current epidemiology, pathogenicity, biofilm formation, natural antifungal products and new therapeutic options. *J. Med. Microbiol.*, **2013**, 62(Pt 1), 10-24. <http://dx.doi.org/10.1099/jmm.0.045054-0> PMID: 23180477
- [3] Arendrup, M.C.; Patterson, T.F. Multidrug-Resistant *Candida*: Epidemiology, Molecular Mechanisms, and Treatment. *J. Infect. Dis.*, **2017**, 216(suppl\_3), S445-S451. <http://dx.doi.org/10.1093/infdis/jix131> PMID: 28911043
- [4] Wisplinghoff, H.; Bischoff, T.; Tallent, S.M.; Seifert, H.; Wenzel, R.P.; Edmond, M.B. Nosocomial bloodstream infections in US hospitals: analysis of 24,179 cases from a prospective nationwide surveillance study. *Clin. Infect. Dis.*, **2004**, 39(3), 309-317. <http://dx.doi.org/10.1086/421946> PMID: 15306996
- [5] Mayer, F.L.; Wilson, D.; Hube, B. *Candida albicans* pathogenicity mechanisms. *Virulence*, **2013**, 4(2), 119-128. <http://dx.doi.org/10.4161/viru.22913> PMID: 23302789
- [6] Tsui, C.; Kong, E.F.; Jabra-Rizk, M.A. Pathogenesis of *Candida albicans* biofilm. *Pathog. Dis.*, **2016**, 74(4)ftw018 <http://dx.doi.org/10.1093/femspd/ftw018> PMID: 26960943
- [7] Whaley, S.G.; Berkow, E.L.; Rybak, J.M.; Nishimoto, A.T.; Barker, K.S.; Rogers, P.D. Azole Antifungal Resistance in *Candida albicans* and Emerging Non-*albicans Candida* Species. *Front. Microbiol.*, **2017**, 7, 2173. <http://dx.doi.org/10.3389/fmicb.2016.02173> PMID: 28127295
- [8] Arif, T.; Bhosale, J.D.; Kumar, N.; Mandal, T.K.; Bendre, R.S.; Lavekar, G.S.; Dabur, R. Natural products-antifungal agents derived from plants. *J. Asian Nat. Prod. Res.*, **2009**, 11(7), 621-638. <http://dx.doi.org/10.1080/10286020902942350> PMID: 20183299
- [9] Vidyasagar, G.M. Plant-Derived Antifungal Agents: Past and Recent Developments. *Recent Trends in Antifungal Agents and Antifungal Therapy*; Basak, Chakraborty, M., Ed.; Springer: India, **2016**, pp. 123-147. [http://dx.doi.org/10.1007/978-81-322-2782-3\\_5](http://dx.doi.org/10.1007/978-81-322-2782-3_5)
- [10] Kier, G.; Kreft, H.; Lee, T.M.; Jetz, W.; Ibsch, P.L.; Nowicki, C.; Mutke, J.; Barthlott, W. A global assessment of endemism and species richness across island and mainland regions. *Proc. Natl. Acad. Sci. USA*, **2009**, 106(23), 9322-9327. <http://dx.doi.org/10.1073/pnas.0810306106> PMID: 19470638
- [11] Yaralizadeh, M.; Abedi, P.; Namjoyan, F.; Fatahinia, M.; Nezamivand Chegini, S. A comparison of the effects of Lawsonia inermis (Iranian henna) and clotrimazole on *Candida albicans* in rats. *J. Mycol. Med.*, **2018**, 28(3), 419-423. <http://dx.doi.org/10.1016/j.mycmed.2018.05.012> PMID: 29891221
- [12] Akroum, S. Antifungal activity of acetone extracts from *Punica granatum* L., *Quercus suber* L. and *Vicia faba* L. *J. Mycol. Med.*, **2017**, 27(1), 83-89. <http://dx.doi.org/10.1016/j.mycmed.2016.10.004> PMID: 27856170
- [13] Akroum, S. Antifungal activity of *Camellia sinensis* crude extracts against four species of *Candida* and *Microsporum persicolor*. *J. Mycol. Med.*, **2018**, 28(3), 424-427. <http://dx.doi.org/10.1016/j.mycmed.2018.06.003> PMID: 29960870
- [14] Favre-Godal, Q.; Dorsaz, S.; Queiroz, E.F.; Conan, C.; Marcourt, L.; Wardojo, B.P.; Voinesco, F.; Buchwalder, A.; Gindro, K.; Sanglard, D.; Wolfender, J.L. Comprehensive approach for the detection of antifungal compounds using a susceptible strain of *Candida albicans* and confirmation of *in vivo* activity with the *Galleria mellonella* model. *Phytochemistry*, **2014**, 105, 68-78. <http://dx.doi.org/10.1016/j.phytochem.2014.06.004> PMID: 24984572
- [15] de Campos Rasteiro, M.M.; da Costa, A.C.; Araújo, C.F.; de Barros, P.P.; Rossoni, R.D. Essential oil of *Melaleuca alternifolia* for the treatment of oral candidiasis induced in an immunosuppressed mouse model. *BMC Complement. Altern. Med.*, **2014**, 14, 489. <http://dx.doi.org/10.1186/1472-6882-14-489> PMID: 25510285
- [16] Campos, L.M.; de Melo, L.; Lemos, A.S.O.; Guedes, M.C.M.R.; Silva, T.P.; Figueiredo, G.F.; Reis, J.L.; Rocha, V.N.; Melo, R.C.N.; Araújo, M.G.F.; Apolônio, A.C.M.; Scio, E.; Fabri, R.L. *Mitracarpus frigidus*: A promising antifungal in the treatment of vulvovaginal candidiasis. *Ind. Crops Prod.*, **2018**, 123, 731-739. <http://dx.doi.org/10.1016/j.indcrop.2018.07.038>
- [17] Bitencourt, P.E.; Cargnelutti, L.O.; Stein, C.S.; Lautenchleger, R.; Ferreira, L.M.; Sangoi, M.; Denardi, L.; Borges, R.M.; Boligon, A.; Moresco, R.N.; Cruz, L.; Zanette, R.A.; Alves, S.H.; Moretto, M.B. Nanoparticle formulation increases *Syzygium cumini* antioxidant activity in *Candida albicans*-infected diabetic rats. *Pharm. Biol.*, **2017**, 55(1), 1082-1088. <http://dx.doi.org/10.1080/13880209.2017.1283338> PMID: 28193098
- [18] Yang, F.; Xiao, C.; Qu, J.; Wang, G. Structural characterization of low molecular weight polysaccharide from *Astragalus membranaceus* and its immunologic enhancement in recombinant protein vaccine against systemic candidiasis. *Carbohydr. Polym.*, **2016**, 145, 48-55. <http://dx.doi.org/10.1016/j.carbpol.2016.03.024> PMID: 27106150
- [19] Rampadarath, S.; Puchoo, D.; Jeewon, R. *Jatropha curcas* L: Phytochemical, antimicrobial and larvicidal properties. *Asian Pac. J. Trop. Biomed.*, **2016**, 6, 858-865. <http://dx.doi.org/10.1016/j.apjtb.2016.01.019>
- [20] Zorić, N.; Kopjar, N.; Kraljić, K.; Oršolić, N.; Tomić, S.; Kosalec, I. Olive leaf extract activity against *Candida albicans* and *C. dubliniensis* - the *in vitro* viability study. *Acta Pharm.*, **2016**, 66(3), 411-421. <http://dx.doi.org/10.1515/acph-2016-0033> PMID: 27383889
- [21] Mertas, A.; Garbusińska, A.; Szliszka, E.; Jureczko, A.; Kowalska, M.; Król, W. The influence of tea tree oil (*Melaleuca alternifolia*) on fluconazole activity against fluconazole-resistant *Candida albicans* strains. *BioMed Res. Int.*, **2015**, 2015590470 <http://dx.doi.org/10.1155/2015/590470> PMID: 25722982
- [22] de Freitas, A.L.D.; Kaplum, V.; Rossi, D.C.P.; da Silva, L.B.R.; Melhem, M.S.C.; Taborda, C.P.; de Mello, J.C.P.; Nakamura, C.V.; Ishida, K. Proanthocyanidin polymeric tannins from *Stryphnodendron adstringens* are effective against *Candida* spp. isolates and for vaginal candidiasis treatment. *J. Ethnopharmacol.*, **2018**, 216, 184-190. <http://dx.doi.org/10.1016/j.jep.2018.01.008> PMID: 29325916
- [23] Bajpai, V.K.; Park, Y.H.; Kang, S.C. A diterpenoid taxodone from *Metasequoia glyptostroboides* with antimycotic potential against clinical isolates of *Candida* species. *J. Mycol. Med.*, **2015**, 25(1), e31-e38. <http://dx.doi.org/10.1016/j.mycmed.2014.10.017> PMID: 25637428
- [24] Correia, A.F.; Silveira, D.; Fonseca-Bazzo, Y.M.; Magalhães, P.O.; Fagg, C.W.; da Silva, E.C.; Gomes, S.M.; Gandolfi, L.; Pratesi, R.; de Medeiros Nóbrega, Y.K. Activity of crude extracts from Brazilian cerrado plants against clinically relevant *Candida* species. *BMC Complement. Altern. Med.*, **2016**, 16, 203. <http://dx.doi.org/10.1186/s12906-016-1164-3> PMID: 27401815
- [25] Mgbeahuruike, E.E.; Holm, Y.; Vuorela, H.; Amandikwa, C.; Fyhrquist, P. An ethnobotanical survey and antifungal activity of *Piper guineense* used for the treatment of fungal infections in West-African traditional medicine. *J. Ethnopharmacol.*, **2019**, 229, 157-166. <http://dx.doi.org/10.1016/j.jep.2018.10.005> PMID: 30336302
- [26] Giongo, J.L.; de Almeida Vaucher, R.; Fausto, V.P.; Quatrin, P.M.; Lopes, L.Q.S.; Santos, R.C.V.; Gündel, A.; Gomes, P.; Steppe, M. Anti-*Candida* activity assessment of *Pelargonium graveolens* oil free and nanoemulsion in biofilm formation in hospital medical supplies. *Microb. Pathog.*, **2016**, 100, 170-178. <http://dx.doi.org/10.1016/j.micpath.2016.08.013> PMID: 27544324
- [27] Favre-Godal, Q.; Dorsaz, S.; Queiroz, E.F.; Marcourt, L.; Ebrahimi, S.N.; Allard, P.M.; Voinesco, F.; Hamburger, M.; Gupta, M.P.;

- Gindro, K.; Sanglard, D.; Wolfender, J.L. Anti- Candida Cassane-Type Diterpenoids from the Root Bark of Swartzia simplex. *J. Nat. Prod.*, **2015**, *78*, 2994-3004.  
<http://dx.doi.org/10.1021/acs.jnatprod.5b00744>
- [28] Rivera-Yañez, C.R.; Terrazas, L.I.; Jimenez-Estrada, M.; Campos, J.E.; Flores-Ortiz, C.M.; Hernandez, L.B.; Cruz-Sanchez, T.; Garrido-Fariña, G.I.; Rodriguez-Monroy, M.A.; Canales-Martinez, M.M. Anti-Candida Activity of Bursera morelensis Ramirez Essential Oil and Two Compounds,  $\alpha$ -Pinene and  $\gamma$ -Terpinene-An *In Vitro* Study. *Molecules*, **2017**, *22*(12), 2095.  
<http://dx.doi.org/10.3390/molecules22122095> PMID: 29206158
- [29] Ngo-Mback, M.N.L.N.L. MubarakAli, D.; Dongmo, P.M.J.M.J.; Boyom, F.F.; Thajuddin, N. Anti-Candidal biofilm potential of solvent extracts of *Aeollanthus cucullatus* (Ryding) and its chemical analysis. *Biocatal. Agric. Biotechnol.*, **2019**, *17*, 595-604.  
<http://dx.doi.org/10.1016/j.cbac.2019.01.012>
- [30] de Morais, C.B.; Scopel, M.; Pedrazza, G.P.R.; da Silva, F.K.; Dalla Lana, D.F.; Tonello, M.L.; Miotto, S.T.S.; Machado, M.M.; De Oliveira, L.F.S.; Fuentesfria, A.M.; Zuanazzi, J.A.S. Antidermatophyte activity of Leguminosae plants from Southern Brazil with emphasis on *Mimosa pigra* (Leguminosae). *J. Mycol. Med.*, **2017**, *27*(4), 530-538.  
<http://dx.doi.org/10.1016/j.mycmed.2017.07.006> PMID: 28822705
- [31] Suurbaar, J.; Mosobil, R.; Donkor, A.M.M. Antibacterial and antifungal activities and phytochemical profile of leaf extract from different extractants of *Ricinus communis* against selected pathogens. *BMC Res. Notes*, **2017**, *10*(1), 660.  
<http://dx.doi.org/10.1186/s13104-017-3001-2> PMID: 29191226
- [32] Carvalho, R.S.; Carollo, C.A.; de Magalhães, J.C.; Palumbo, J.M.C.; Boaretto, A.G.; Nunes e Sá, I.C.; Ferraz, A.C.; Lima, W.G.; de Siqueira, J.M.; Ferreira, J.M.S. Antibacterial and antifungal activities of phenolic compound-enriched ethyl acetate fraction from *Cochlospermum regium* (mart. Et. Schr.) Pilger roots: Mechanisms of action and synergism with tannin and gallic acid. *S. Afr. J. Bot.*, **2018**, *114*, 181-187.  
<http://dx.doi.org/10.1016/j.sajb.2017.11.010>
- [33] Bisio, A.; Schito, A.M.; Ebrahimi, S.N.; Hamburger, M.; Mele, G.; Piatti, G.; Romussi, G.; Dal Piaz, F.; De Tommasi, N. Antibacterial compounds from *Salvia adenophora* Fernald (Lamiaceae). *Phytochemistry*, **2015**, *110*, 120-132.  
<http://dx.doi.org/10.1016/j.phytochem.2014.10.033> PMID: 25435172
- [34] Masoko, P.; Makgapeetja, D.M. Antibacterial, antifungal and antioxidant activity of *Olea africana* against pathogenic yeast and nosocomial pathogens. *BMC Complement. Altern. Med.*, **2015**, *15*, 409.  
<http://dx.doi.org/10.1186/s12906-015-0941-8> PMID: 26577343
- [35] Kutluk, I.; Aslan, M.; Orhan, I.E.E.; Özçelik, B. Antibacterial, antifungal and antiviral bioactivities of selected *Helichrysum* species. *S. Afr. J. Bot.*, **2018**, *119*, 252-257.  
<http://dx.doi.org/10.1016/j.sajb.2018.09.009>
- [36] Seebaluck-Sandoram, R.; Lall, N.; Fibrich, B.; Blom van Staden, A.; Mahomoodally, F. Antibiotic-potential, antioxidant, cytotoxic, anti-inflammatory and anti-acetylcholinesterase potential of *Antidesma madagascariense* Lam. (Euphorbiaceae). *S. Afr. J. Bot.*, **2017**, *111*, 194-201.  
<http://dx.doi.org/10.1016/j.sajb.2017.03.034>
- [37] Minooeianhaghghi, M.H.; Sepehrian, L.; Shokri, H. Antifungal effects of *Lavandula binaludensis* and *Cuminum cyminum* essential oils against *Candida albicans* strains isolated from patients with recurrent vulvovaginal candidiasis. *J. Mycol. Med.*, **2017**, *27*(1), 65-71.  
<http://dx.doi.org/10.1016/j.mycmed.2016.09.002> PMID: 27751723
- [38] Samadi, F.M.; Suhail, S.; Sonam, M.; Sharma, N.; Singh, S.; Gupta, S.; Dobhal, A.; Pradhan, H. Antifungal efficacy of herbs. *J. Oral Biol. Craniofac. Res.*, **2019**, *9*(1), 28-32.  
<http://dx.doi.org/10.1016/j.jobcr.2018.06.002> PMID: 30197861
- [39] Essid, R.; Hammami, M.; Gharbi, D.; Karkouch, I.; Hamouda, T.B.; Elkahoui, S.; Limam, F.; Tabbene, O. Antifungal mechanism of the combination of *Cinnamomum verum* and *Pelargonium graveolens* essential oils with fluconazole against pathogenic *Candida* strains. *Appl. Microbiol. Biotechnol.*, **2017**, *101*(18), 6993-7006.  
<http://dx.doi.org/10.1007/s00253-017-8442-y> PMID: 28766033
- [40] Pereira, J.V.; Freires, I.A.; Castilho, A.R.; da Cunha, M.G.; Alves, Hda.S.; Rosalen, P.L. Antifungal potential of *Sideroxylon obtusifolium* and *Syzygium cumini* and their mode of action against *Candida albicans*. *Pharm. Biol.*, **2016**, *54*(10), 2312-2319.  
<http://dx.doi.org/10.3109/13880209.2016.1155629> PMID: 26987037
- [41] Njateng, G.S.; Du, Z.; Gatsing, D.; Nanfack Donfack, A.R.; Feussi Talla, M.; Kamdem Wabo, H.; Tane, P.; Mouokeu, R.S.; Luo, X.; Kuate, J.R. Antifungal properties of a new terpenoid saponin and other compounds from the stem bark of *Polyscias fulva* Hiern (Araliaceae). *BMC Complement. Altern. Med.*, **2015**, *15*, 25.  
<http://dx.doi.org/10.1186/s12906-015-0541-7> PMID: 25880310
- [42] Yessoufou, K.; Elansary, H.O.; Mahmoud, E.A.; Skalicka-Woźniak, K. Antifungal, antibacterial and anticancer activities of *Ficus drupacea* L. stem bark extract and biologically active isolated compounds. *Ind. Crops Prod.*, **2015**, *74*, 752-758.  
<http://dx.doi.org/10.1016/j.indcrop.2015.06.011>
- [43] Boulaaba, M.; Snoussi, M.; Saada, M.; Mkdmini, K.; Smaoui, A.; Abdelly, C.; Ksouri, R. Antimicrobial activities and phytochemical analysis of *Tamarix gallica* extracts. *Ind. Crops Prod.*, **2015**, *76*, 1114-1122.  
<http://dx.doi.org/10.1016/j.indcrop.2015.08.020>
- [44] Ahmed, D.; Saeed, R.; Shakeel, N.; Fatima, K.; Arshad, A. Antimicrobial activities of methanolic extract of *Carissa opaca* roots and its fractions and compounds isolated from the most active ethyl acetate fraction. *Asian Pac. J. Trop. Biomed.*, **2015**, *5*, 541-545.  
<http://dx.doi.org/10.1016/j.apjtb.2015.05.006>
- [45] Juliano, C.; Marchetti, M.; Campagna, P.; Usai, M. Antimicrobial activity and chemical composition of essential oil from *Helichrysum microphyllum* Cambess. subsp. *tyrrhenicum* Bacch., *Bruello & Giusso* collected in South-West Sardinia. *Saudi J. Biol. Sci.*, **2018**, ...  
<http://dx.doi.org/10.1016/j.sjbs.2018.04.009> PMID: 31303817
- [46] Costa, A.R.; de Lima Silva, J.; Lima, K.R.R.; Rocha, M.I.; Barros, L.M.; da Costa, J.G.M.; Boligon, A.A.; Kamdem, J.P.; Carneiro, J.N.P.; Leite, N.F.; de Menezes, I.R.A.; Duarte, A.E.; Morais-Braga, M.F.B.; Coutinho, H.D.M. *Rhaphiodon echinus* (Nees & Mart.) Schauer: Chemical, toxicological activity and increased antibiotic activity of antifungal drug activity and antibacterial. *Microb. Pathog.*, **2017**, *107*, 280-286.  
<http://dx.doi.org/10.1016/j.micpath.2017.04.001> PMID: 28385578
- [47] Bottari, N.B.; Lopes, L.Q.; Pizzuti, K.; Filippi Dos Santos Alves, C.; Corrêa, M.S.; Bolzan, L.P.; Zago, A.; de Almeida Vaucher, R.; Boligon, A.A.; Giongo, J.L.; Baldissera, M.D.; Santos, R.C. Antimicrobial activity and phytochemical characterization of *Carya illinoensis*. *Microb. Pathog.*, **2017**, *104*, 190-195.  
<http://dx.doi.org/10.1016/j.micpath.2017.01.037> PMID: 28126664
- [48] Cavalcanti Filho, J.R.; Silva, T.F.; Nobre, W.Q.; Oliveira de Souza, L.I.; Silva E Silva Figueiredo, C.S.; Figueiredo, R.C.; de Gusmão, N.B.; Silva, M.V.; Nascimento da Silva, L.C.; Correia, M.T. Antimicrobial activity of *Buchenavia tetraphylla* against *Candida albicans* strains isolated from vaginal secretions. *Pharm. Biol.*, **2017**, *55*(1), 1521-1527.  
<http://dx.doi.org/10.1080/13880209.2017.1304427> PMID: 28376640
- [49] Packer, J.; Naz, T.; Harrington, D.; Jamie, J.F.; Vemulpad, S.R. Yaegl Community Elders. Antimicrobial activity of customary medicinal plants of the Yaegl Aboriginal community of northern New South Wales, Australia: a preliminary study. *BMC Res. Notes*, **2015**, *8*, 276.  
<http://dx.doi.org/10.1186/s13104-015-1258-x> PMID: 26122212
- [50] Ebani, V.V.; Nardon, S.; Bertelloni, F.; Pistelli, L.; Mancianti, F. Antimicrobial activity of five essential oils against bacteria and fungi responsible for urinary tract infections. *Molecules*, **2018**, *23*(7), 1-12.  
<http://dx.doi.org/10.3390/molecules23071668> PMID: 29987237
- [51] Nair, S.V.; Baranwal, G.; Chatterjee, M.; Sachu, A.; Vasudevan, A.K.; Bose, C.; Banerji, A.; Biswas, R. Antimicrobial activity of plumbagin, a naturally occurring naphthoquinone from *Plumbago rosea*, against *Staphylococcus aureus* and *Candida albicans*. *Int. J. Med. Microbiol.*, **2016**, *306*(4), 237-248.  
<http://dx.doi.org/10.1016/j.ijmm.2016.05.004> PMID: 27212459



- [52] Besbes Hlila, M.; Mosbah, H.; Majouli, K.; Ben Nejma, A.; Ben Jannet, H.; Mastouri, M.; Aouni, M.; Selmi, B. Antimicrobial Activity of *Scabiosa arenaria* Forssk. Extracts and Pure Compounds Using Bioguided Fractionation. *Chem. Biodivers.*, **2016**, *13*(10), 1262-1272.  
<http://dx.doi.org/10.1002/cbdv.201600028> PMID: 27448449
- [53] Lulekal, E.; Rondevaldova, J.; Bernaskova, E.; Cepkova, J.; Asfaw, Z.; Kelbessa, E. Antimicrobial activity of traditional medicinal plants from Ankober District, North Shewa Zone, Amhara Region, Ethiopia. *Pharm. Biol.*, **2014**, *52*(5), 614-620.  
<http://dx.doi.org/10.3109/13880209.2013.858362> PMID: 24392738
- [54] Hlila, M.B.; Majouli, K.; Jannet, H.B.; Aouni, M.; Mastouri, M.; Selmi, B. Antimicrobial activity of Tunisian *Euphorbia paralias* L. *Asian Pac. J. Trop. Biomed.*, **2017**, *7*, 629-632.  
<http://dx.doi.org/10.1016/j.apjtb.2017.06.008>
- [55] Quatrin, P.M.; Verdi, C.M.; de Souza, M.E.; de Godoi, S.N.; Klein, B.; Gundel, A.; Wagner, R.; de Almeida Vaucher, R.; Ourique, A.F.; Santos, R.C.V. Antimicrobial and antibiofilm activities of nanoemulsions containing *Eucalyptus globulus* oil against *Pseudomonas aeruginosa* and *Candida* spp. *Microb. Pathog.*, **2017**, *112*, 230-242.  
<http://dx.doi.org/10.1016/j.micpath.2017.09.062> PMID: 28970174
- [56] Dube, P.; Meyer, S.; Marnewick, J.L.L. Antimicrobial and antioxidant activities of different solvent extracts from fermented and green honeybush (*Cyclopia intermedia*) plant material. *S. Afr. J. Bot.*, **2017**, *110*, 184-193.  
<http://dx.doi.org/10.1016/j.sajb.2016.10.010>
- [57] Akter, K.; Barnes, E.C.; Loa-Kum-Cheung, W.L.; Yin, P.; Kichu, M.; Brophy, J.J.; Barrow, R.A.; Imchen, I.; Vemulpad, S.R.; Jamie, J.F. Antimicrobial and antioxidant activity and chemical characterisation of *Erythrina stricta* Roxb. (Fabaceae). *J. Ethnopharmacol.*, **2016**, *185*, 171-181.  
<http://dx.doi.org/10.1016/j.jep.2016.03.011> PMID: 26969405
- [58] Göger, G.; Demirci, B.; Ilgin, S.; Demirci, F. Antimicrobial and toxicity profiles evaluation of the Chamomile (*Matricaria recutita* L.) essential oil combination with standard antimicrobial agents. *Ind. Crops Prod.*, **2018**, *120*, 279-285.  
<http://dx.doi.org/10.1016/j.indcrop.2018.04.024>
- [59] Costa, G.M.; Endo, E.H.; Cortez, D.A.; Nakamura, T.U.; Nakamura, C.V.; Dias Filho, B.P. Antimicrobial effects of *Piper hispidum* extract, fractions and chalcones against *Candida albicans* and *Staphylococcus aureus*. *J. Mycol. Med.*, **2016**, *26*(3), 217-226.  
<http://dx.doi.org/10.1016/j.mycmed.2016.03.002> PMID: 27499460
- [60] Emmanuel, R.; Saravanan, M.; Ovais, M.; Padmavathy, S.; Shinwari, Z.K.; Prakash, P. Antimicrobial efficacy of drug blended biosynthesized colloidal gold nanoparticles from *Justicia glauca* against oral pathogens: A nanoantibiotic approach. *Microb. Pathog.*, **2017**, *113*, 295-302.  
<http://dx.doi.org/10.1016/j.micpath.2017.10.055> PMID: 29101061
- [61] Ramadwa, T.E.; Elgorashi, E.E.; McGaw, L.J.; Ahmed, A.S.; Eloff, J.N. Antimicrobial, anti-inflammatory activity and cytotoxicity of *Funtumia africana* leaf extracts, fractions and the isolated methyl ursolate. *S. Afr. J. Bot.*, **2017**, *108*, 126-131.  
<http://dx.doi.org/10.1016/j.sajb.2016.10.019>
- [62] Pendota, S.C.; Aderogba, M.A.; Moyo, M.; McGaw, L.J.; Mulaudzi, R.B.; Van Staden, J. Antimicrobial, antioxidant and cytotoxicity of isolated compounds from leaves of *Pappea capensis*. *S. Afr. J. Bot.*, **2017**, *108*, 272-277.  
<http://dx.doi.org/10.1016/j.sajb.2016.10.021>
- [63] Dos Santos Alves, C.F.; Bonez, P.C.; de Souza, M.E.; da Cruz, R.C.; Boligon, A.A.; Piana, M.; Brum, T.F.; Rossi, G.G.; Jesus, R.D.; Grando, T.H.; Monteiro, S.G.; Anraku de Campos, M.M.; Giongo, J.L.; Vianna Santos, R.C. Antimicrobial, antitypanosomal and antibiofilm activity of *Equisetum hyemale*. *Microb. Pathog.*, **2016**, *101*, 119-125.  
<http://dx.doi.org/10.1016/j.micpath.2016.11.008> PMID: 27856271
- [64] Leite, T.R.; Silva, M.A.P.D.; Santos, A.C.B.D.; Coutinho, H.D.M.; Duarte, A.E.; Costa, J.G.M.D. Antimicrobial, modulatory and chemical analysis of the oil of *Croton limae*. *Pharm. Biol.*, **2017**, *55*(1), 2015-2019.  
<http://dx.doi.org/10.1080/13880209.2017.1355926> PMID: 28738715
- [65] Ruiz-Pérez, N.J.; González-Ávila, M.; Sánchez-Navarrete, J.; Toscano-Garibay, J.D. Antimycotic Activity and Genotoxic Evaluation of *Citrus sinensis* and *Citrus latifolia* Essential Oils. *Sci. Rep.*, **2016**, *6*, 25371.  
<http://dx.doi.org/10.1038/srep25371> PMID: 27137128
- [66] Hamdi, A.; Majouli, K.; Flamini, G.; Marzouk, B.; Marzouk, Z.; Heyden, Y.V. Antioxidant and anticandidal activities of the Tunisian *Haplophyllum tuberculatum* (Forssk.) A. Juss. essential oils. *S. Afr. J. Bot.*, **2017**, *112*, 210-214.  
<http://dx.doi.org/10.1016/j.sajb.2017.05.026>
- [67] Camargo, L.E.; Pedroso, L.S.; Vendrame, S.C.; Mainardes, R.M.; Khalil, N.M. Antioxidant and antifungal activities of *Camellia sinensis* (L.) Kuntze leaves obtained by different forms of production. *Braz. J. Biol.*, **2016**, *76*(2), 428-434.  
<http://dx.doi.org/10.1590/1519-6984.18814> PMID: 26983085
- [68] Abu Zarin, M.; Wan, H.Y.; Isha, A.; Armania, N. Antioxidant, antimicrobial and cytotoxic potential of condensed tannins from *Leucaena leucocephala* hybrid-Rendang. *Food Sci. Hum. Wellness*, **2016**, *5*, 65-75.  
<http://dx.doi.org/10.1016/j.fshw.2016.02.001>
- [69] Knežević, A.; Stajić, M.; Sofrenić, I.; Stanojković, T.; Milovanović, I.; Tešević, V. Antioxidative, antifungal, cytotoxic and antineurodegenerative activity of selected *Trametes* species from Serbia. *PLoS One*, **2018**, *13*(8), e0203064  
<http://dx.doi.org/10.1371/journal.pone.0203064> PMID: 30169542
- [70] Abu-Darwish, M.S.; Cabral, C.; Gonçalves, M.J.; Cavaleiro, C.; Cruz, M.T.; Effert, T.; Salgueiro, L. *Artemisia herba-alba* essential oil from Buseirah (South Jordan): Chemical characterization and assessment of safe antifungal and anti-inflammatory doses. *J. Ethnopharmacol.*, **2015**, *174*, 153-160.  
<http://dx.doi.org/10.1016/j.jep.2015.08.005> PMID: 26277492
- [71] Soliman, S.S.M.; Semreen, M.H.; El-Keblawy, A.A.; Abdullah, A.; Uppuluri, P.; Ibrahim, A.S. Assessment of herbal drugs for promising anti-*Candida* activity. *BMC Complement. Altern. Med.*, **2017**, *17*(1), 257.  
<http://dx.doi.org/10.1186/s12906-017-1760-x> PMID: 28482836
- [72] Lee, W.; Woo, E.R.; Lee, D.G. Effect of apigenin isolated from *Aster yomena* against *Candida albicans*: apigenin-triggered apoptotic pathway regulated by mitochondrial calcium signaling. *J. Ethnopharmacol.*, **2019**, *231*, 19-28.  
<http://dx.doi.org/10.1016/j.jep.2018.11.005> PMID: 30408533
- [73] Linhares Neto, M.V.; da Silva, R.O.; de Oliveira, F.F.; Costa, L.C.B.; Conceição, A.O.; de Oliveira, R.A. Avaliação anti-*Candida* of essential oils from three medicinal plants species (Asteraceae). *S. Afr. J. Bot.*, **2018**, *115*, 132-137.  
<http://dx.doi.org/10.1016/j.sajb.2018.01.017>
- [74] Buhian, W.P.C.; Rubio, R.O.; Valle, D.L.; Martin-Puzon, J.J. Bioactive metabolite profiles and antimicrobial activity of ethanolic extracts from *Muntingia calabura* L. leaves and stems. *Asian Pac. J. Trop. Biomed.*, **2016**, *6*, 682-685.  
<http://dx.doi.org/10.1016/j.apjtb.2016.06.006>
- [75] Panyo, J.; Matsunami, K.; Panichayupakaranant, P. Bioassay-guided isolation and evaluation of antimicrobial compounds from *Ixora megalophylla* against some oral pathogens. *Pharm. Biol.*, **2016**, *54*(9), 1522-1527.  
<http://dx.doi.org/10.3109/13880209.2015.1107106> PMID: 26809027
- [76] Yang, Y.; Chen, H.; Lei, J.; Yu, J. Biological activity of extracts and active compounds isolated from *Siegesbeckia orientalis* L. *Ind. Crops Prod.*, **2016**, *94*, 288-293.  
<http://dx.doi.org/10.1016/j.indcrop.2016.08.023>
- [77] Malik, T.A.; Kamili, A.N.; Chishti, M.Z.; Ahad, S.; Tantry, M.A.; Hussain, P.R.; Johri, R.K. Breaking the resistance of *Escherichia coli*: Antimicrobial activity of *Berberis lycium* Royle. *Microb. Pathog.*, **2017**, *102*, 12-20.  
<http://dx.doi.org/10.1016/j.micpath.2016.11.011> PMID: 27888048
- [78] Borah, B.; Phukon, P.; Hazarika, M.; Ahmed, R.; Sarmah, D.; Wann, S.B.; Das, A.; Bhau, B. *Calamus leptospadix* Griff. a high saponin yielding plant with antimicrobial property. *Ind. Crops Prod.*, **2016**, *82*, 127-132.  
<http://dx.doi.org/10.1016/j.indcrop.2015.11.075>
- [79] Shinobu-Mesquita, C.S.; Bonfim-Mendonça, P.S.; Moreira, A.L.; Ferreira, I.C.; Donatti, L.; Fiorini, A.; Svidzinski, T.I. Cellular

- Structural Changes in *Candida albicans* Caused by the Hydroalcoholic Extract from *Sapindus saponaria* L. *Molecules*, **2015**, *20*(5), 9405-9418.  
<http://dx.doi.org/10.3390/molecules20059405> PMID: 26007191
- [80] Padalia, H.; Chanda, S. Characterization, antifungal and cytotoxic evaluation of green synthesized zinc oxide nanoparticles using *Ziziphus nummularia* leaf extract. *Artif. Cells Nanomed. Biotechnol.*, **2017**, *45*(8), 1751-1761.  
<http://dx.doi.org/10.1080/21691401.2017.1282868> PMID: 28140658
- [81] Falcão, S.; Bacém, I.; Igrejas, G.; Rodrigues, P.J.; Vilas-Boas, M.; Joana, S. Chemical composition and antimicrobial activity of hydrodistilled oil from juniper berries. *Ind. Crops Prod.*, **2018**, *124*, 878-884.  
<http://dx.doi.org/10.1016/j.indcrop.2018.08.069>
- [82] Abu-Darwish, M.S.; Cabral, C.; Gonçalves, M.J.; Cavaleiro, C.; Cruz, M.T.; Zulfıqar, A.; Khan, I.A.; Efferth, T.; Salgueiro, L. Chemical composition and biological activities of *Artemisia judaica* essential oil from southern desert of Jordan. *J. Ethnopharmacol.*, **2016**, *191*, 161-168.  
<http://dx.doi.org/10.1016/j.jep.2016.06.023> PMID: 27318275
- [83] Batiga, S.; Valli, M.; Zeraik, M.L.; Fraige, K.; Leme, G.M.; Pitanqui, N.S.; Almeida, A.M.F.; Michel, S.; Young, M.C.M.; Bolzani, V.S. Chemical composition and biological properties of *Ipomoea procumbens*. *Braz. J. Pharmacognosy*, **2018**, *29*, 191-197.  
<http://dx.doi.org/10.1016/j.bjpp.2018.08.010>
- [84] Dos Santos, J.F.S.; Rocha, J.E.; Bezerra, C.F.; do Nascimento Silva, M.K.; de Matos, Y.M.L.S.; de Freitas, T.S.; Dos Santos, A.T.L.; da Cruz, R.P.; Machado, A.J.T.; Rodrigues, T.H.S.; de Brito, E.S.; Sales, D.L.; de Oliveira Almeida, W.; da Costa, J.G.M.; Coutinho, H.D.M.; Morais-Braga, M.F.B. Chemical composition, antifungal activity and potential anti-virulence evaluation of the *Eugenia uniflora* essential oil against *Candida* spp. *Food Chem.*, **2018**, *261*, 233-239.  
<http://dx.doi.org/10.1016/j.foodchem.2018.04.015> PMID: 29739588
- [85] Khosravi, A.R.; Sharifzadeh, A.; Nikaein, D.; Almaie, Z.; Gandomi Nasrabadi, H. Chemical composition, antioxidant activity and antifungal effects of five Iranian essential oils against *Candida* strains isolated from urine samples. *J. Mycol. Med.*, **2018**, *28*(2), 355-360.  
<http://dx.doi.org/10.1016/j.mycmed.2018.01.005> PMID: 29477783
- [86] Nogueira Sobrinho, A.C.; Bezerra de Souza, E.; Rocha, M.F.G.; Ribeiro Albuquerque, M.R.J.; Nogueira Bandeira, P.; Silva dos Santos, H.; de Paula Cavalcante, C.S.; Souza Oliveira, S.; Rodrigues Aragão, P.; Maia de Moraes, S.; Oliveira dos Santos Fontenelle, R. Chemical composition, antioxidant, antifungal and hemolytic activities of essential oil from *Baccharis trinervis* (Lam.) Pers. (Asteraceae). *Ind. Crops Prod.*, **2016**, *84*, 108-115.  
<http://dx.doi.org/10.1016/j.indcrop.2016.01.051>
- [87] Ertaş, A.; Boğa, M.; Yılmaz, M.A.; Yeşil, Y.; Haşimi, N.; Kaya, M.Ş.; Temel, H.; Kolak, U. Chemical compositions by using LC-MS/MS and GC-MS and biological activities of *Sedum sediforme* (Jacq.) Pau. *J. Agric. Food Chem.*, **2014**, *62*(20), 4601-4609.  
<http://dx.doi.org/10.1021/jf500067q> PMID: 24773044
- [88] Desam, N.R.; Al-Rajab, A.J.; Sharma, M.; Mylabathula, M.M.; Gowkanapalli, R.R.; Albratty, M. Chemical constituents, *in vitro* antibacterial and antifungal activity of *Mentha piperita* L. (peppermint) essential oils *Journal of King Saud University - Science*, **2019**, *31*, 528-533.
- [89] Akarchariya, N.; Sirilun, S.; Julsrigraval, J.; Chansakaowa, S. Chemical profiling and antimicrobial activity of essential oil from *Curcuma aeruginosa* Roxb., *Curcuma glans* K. Larsen & J. Mood and *Curcuma cf. xanthorrhiza* Roxb. collected in Thailand. *Asian Pac. J. Trop. Biomed.*, **2017**, *7*, 881-885.  
<http://dx.doi.org/10.1016/j.apjtb.2017.09.009>
- [90] Gavanji, S.; Larki, B. Comparative effect of propolis of honey bee and some herbal extracts on *Candida albicans*. *Chin. J. Integr. Med.*, **2017**, *23*(3), 201-207.  
<http://dx.doi.org/10.1007/s11655-015-2074-9> PMID: 26149083
- [91] Naeini, A.; Naderi, N.J.; Shokri, H. Analysis and *in vitro* anti-*Candida* antifungal activity of *Cuminum cuminum* and *Salvadora persica* herbs extracts against pathogenic *Candida* strains. *J. Mycol. Med.*, **2014**, *24*(1), 13-18.  
<http://dx.doi.org/10.1016/j.mycmed.2013.09.006> PMID: 24210587
- [92] Lee, H.S.; Kim, Y. Development of *Candida albicans* biofilms is diminished by *Paeonia lactiflora* *via* obstruction of cell adhesion and cell lysis. *J. Microbiol. Biotechnol.*, **2018**, *28*(3), 482-490.  
<http://dx.doi.org/10.4014/jmb.1712.12041> PMID: 29316739
- [93] Li, J.X.; Li, Q.J.; Guan, Y.F.; Song, X.; Liu, Y.H.; Zhang, J.J.; Li, W.F.; Du, J.; Zhu, M.; Banas, J.A.; Li, X.N.; Pan, L.T.; Zhang, H.J. Discovery of antifungal constituents from the Miao medicinal plant *Isodon flavidus*. *J. Ethnopharmacol.*, **2016**, *191*, 372-378.  
<http://dx.doi.org/10.1016/j.jep.2016.06.046> PMID: 27340103
- [94] Klewicka, E.; Sójka, M.; Klewicki, R.; Kołodziejczyk, K.; Lipińska, L.; Nowak, A. Ellagitannins from Raspberry (*Rubus idaeus* L.) Fruit as Natural Inhibitors of *Geotrichum candidum*. *Molecules*, **2016**, *21*(7), 908.  
<http://dx.doi.org/10.3390/molecules21070908> PMID: 27420041
- [95] Farisa Banu, S.; Rubini, D.; Shanmugavelan, P.; Murugan, R.; Gowrishankar, S.; Karutha Pandian, S.; Nithyanand, P. Effects of patchouli and cinnamon essential oils on biofilm and hyphae formation by *Candida* species. *J. Mycol. Med.*, **2018**, *28*(2), 332-339.  
<http://dx.doi.org/10.1016/j.mycmed.2018.02.012> PMID: 29571979
- [96] El Zawawy, N.A.; Hafez, E.E. Efficacy of *Pluchea dioscoridis* leaf extract against pathogenic *Candida albicans*. *J. Infect. Dev. Ctries.*, **2017**, *11*(4), 334-342.  
<http://dx.doi.org/10.3855/jidc.8447> PMID: 28459225
- [97] Yeganegi, M.; Tabatabaei Yazdi, F.; Mortazavi, S.A.; Asili, J.; Alizadeh Behbahani, B. *Equisetum telmateia* extracts: Chemical compositions, antioxidant activity and antimicrobial effect on the growth of some pathogenic strain causing poisoning and infection. *Microb. Pathog.*, **2018**, *116*, 62-67.  
<http://dx.doi.org/10.1016/j.micpath.2018.01.014> PMID: 29331369
- [98] Zhu, H.; Zhou, Q.M.; Peng, C.; Chen, M.H.; Li, X.N.; Lin, D.S.; Xiong, L. Pocahemiketals A and B, two new hemiketals with unprecedented sesquiterpenoid skeletons from *Pogostemon cablin*. *Fitoterapia*, **2017**, *120*, 67-71.  
<http://dx.doi.org/10.1016/j.fitote.2017.05.013> PMID: 28576720
- [99] Witkowska-Banaszczak, E.; Długaszewska, J. Essential oils and hydrophilic extracts from the leaves and flowers of *Succisa pratensis* Moench. and their biological activity. *J. Pharm. Pharmacol.*, **2017**, *69*(11), 1531-1539.  
<http://dx.doi.org/10.1111/jphp.12784> PMID: 28744918
- [100] Li, S.; Shi, H.; Chang, W.; Li, Y.; Zhang, M.; Qiao, Y.; Lou, H. Eudesmane sesquiterpenes from Chinese liverwort are substrates of Cdrs and display antifungal activity by targeting Erg6 and Erg11 of *Candida albicans*. *Bioorg. Med. Chem.*, **2017**, *25*(20), 5764-5771.  
<http://dx.doi.org/10.1016/j.bmc.2017.09.001> PMID: 28935182
- [101] Eidi, S.; Azadi, H.G.; Rahbar, N.; Mehmannaavaz, H.R. Evaluation of antifungal activity of hydroalcoholic extracts of *Citrullus colocynthis* fruit. *J. Herb. Med.*, **2015**, *5*, 36-40.  
<http://dx.doi.org/10.1016/j.hermed.2015.01.003>
- [102] Salari, S.; Bakhshi, T.; Sharififar, F.; Naseri, A.; Ghasemi Nejad Almani, P. Evaluation of antifungal activity of standardized extract of *Salvia rhytidea* Benth. (Lamiaceae) against various *Candida* isolates. *J. Mycol. Med.*, **2016**, *26*(4), 323-330.  
<http://dx.doi.org/10.1016/j.mycmed.2016.06.003> PMID: 27499461
- [103] Alizadeh Behbahani, B.; Imani Fooladi, A.A. Evaluation of phytochemical analysis and antimicrobial activities *Allium* essential oil against the growth of some microbial pathogens. *Microb. Pathog.*, **2018**, *114*, 299-303.  
<http://dx.doi.org/10.1016/j.micpath.2017.11.055> PMID: 29196170
- [104] Popović, V.; Stojković, D.; Nikolić, M.; Heyerick, A.; Petrović, S.; Soković, M.; Niketić, M. Extracts of three *Laserpitium* L. species and their principal components laserpitine and sesquiterpene lactones inhibit microbial growth and biofilm formation by oral *Candida* isolates. *Food Funct.*, **2015**, *6*(4), 1205-1211.  
<http://dx.doi.org/10.1039/C5FO00066A> PMID: 25720441
- [105] Bassyouni, R.H.; Wali, I.E.; Kamel, Z.; Kassim, M.F. Fennel oil: A promising antifungal agent against biofilm forming fluconazole resistant *Candida albicans* causing vulvovaginal candidiasis. *J. Herb. Med.*, **2018**, *15100227*  
<http://dx.doi.org/10.1016/j.hermed.2018.08.002>
- [106] Santos, S.S.; Lorenzoni, A.; Pegoraro, N.S.; Denardi, L.B.; Alves, S.H.; Schaffazick, S.R.; Cruz, L. Formulation and *in vitro* evaluation of coconut oil-core cationic nanocapsules intended for vaginal

- delivery of clotrimazole. *Colloids Surf. B Biointerfaces*, **2014**, *116*, 270-276.  
<http://dx.doi.org/10.1016/j.colsurf.2014.01.011> PMID: 24503350
- [107] Mocan, A.; Zengin, G.; Simirgiotis, M.; Schafberg, M.; Mollica, A.; Vodnar, D.C.; Crişan, G.; Rohn, S. Functional constituents of wild and cultivated Goji (*L. barbarum* L.) leaves: phytochemical characterization, biological profile, and computational studies. *J. Enzyme Inhib. Med. Chem.*, **2017**, *32*(1), 153-168.  
<http://dx.doi.org/10.1080/14756366.2016.1243535> PMID: 28095717
- [108] Jackson, D.N.; Yang, L.; Wu, S.; Kennelly, E.J.; Lipke, P.N. Garcinia xanthochymus Benzophenones Promote Hyphal Apoptosis and Potentiate Activity of Fluconazole against *Candida albicans* Biofilms. *Antimicrob. Agents Chemother.*, **2015**, *59*(10), 6032-6038.  
<http://dx.doi.org/10.1128/AAC.00820-15> PMID: 26195512
- [109] da Costa Cordeiro, B.M.P.; de Lima Santos, N.D.; Ferreira, M.R.A.; de Araújo, L.C.C.; Junior, A.R.C.; da Conceição Santos, A.D.; de Oliveira, A.P.; da Silva, A.G.; da Silva Falcão, E.P.; Dos Santos Correia, M.T.; da Silva Almeida, J.R.G.; da Silva, L.C.N.; Soares, L.A.L.; Napoleão, T.H.; da Silva, M.V.; Paiva, P.M.G. Hexane extract from *Spondias tuberosa* (Anacardiaceae) leaves has antioxidant activity and is an anti-*Candida* agent by causing mitochondrial and lysosomal damages. *BMC Complement. Altern. Med.*, **2018**, *18*(1), 284.  
<http://dx.doi.org/10.1186/s12906-018-2350-2> PMID: 30340567
- [110] Mashjoor, S.; Yousefzadi, M. Holothurians antifungal and antibacterial activity to human pathogens in the Persian Gulf. *J. Mycol. Med.*, **2017**, *27*(1), 46-56.  
<http://dx.doi.org/10.1016/j.mycmed.2016.08.008> PMID: 27641487
- [111] Martins, N.; Ferreira, I.C.F.R.; Henriques, M.; Silva, S. *In vitro* anti-*Candida* activity of *Glycyrrhiza glabra* L. *Ind. Crops Prod.*, **2016**, *83*, 81-85.  
<http://dx.doi.org/10.1016/j.indcrop.2015.12.029>
- [112] Oliveira, F.A.; Rorato, V.C.; Almeida-Apolonio, A.A.; Rodrigues, A.B.; Barros, A.L.; Sangalli, A.; Arena, A.C.; Mota, J.S.; Grisolia, A.B.; Oliveira, K.M.P. *In vitro* antifungal activity of *Myracrodruon urundeuva* Allemão against human vaginal *Candida* species. *An. Acad. Bras. Cienc.*, **2017**, *89*(3)(Suppl.), 2423-2432.  
<http://dx.doi.org/10.1590/0001-3765201720170254> PMID: 28746624
- [113] Mbosso Teinkela, J.E.; Siwe Noundou, X.; Fannang, S.; Meyer, F.; Vardamides, J.C.; Mpondo Mpondo, E.; Krause, R.W.M.; Azebaze, A.G.B.; Nguedia, J.C.A. *In vitro* antimicrobial activity of the methanol extract and compounds from the wood of *Ficus elastica* Roxb. ex Hornem. aerial roots. *S. Afr. J. Bot.*, **2017**, *111*, 302-306.  
<http://dx.doi.org/10.1016/j.sajb.2017.03.026>
- [114] Snene, A.; El Mokni, R.; Jmii, H.; Jlassi, I.; Jaïdane, H.; Falconieri, D.; Piras, A.; Dhaouadi, H.; Porcedda, S.; Hammami, S. *In vitro* antimicrobial, antioxidant and antiviral activities of the essential oil and various extracts of wild (*Daucus virgatus* (Poir.) Maire) from Tunisia. *Ind. Crops Prod.*, **2017**, *109*, 109-115.  
<http://dx.doi.org/10.1016/j.indcrop.2017.08.015>
- [115] D'Arrigo, M.; Bisignano, C.; Irrera, P.; Smeriglio, A.; Zagami, R.; Trombetta, D.; Romeo, O.; Mandalari, G. *In vitro* evaluation of the activity of an essential oil from *Pistacia vera* L. variety Bronte hull against *Candida* sp. *BMC Complement. Altern. Med.*, **2019**, *19*(1), 6.  
<http://dx.doi.org/10.1186/s12906-018-2425-0> PMID: 30612544
- [116] Onivogui, G.; Letsididi, R.; Diaby, M.; Wang, L.; Song, Y. Influence of extraction solvents on antioxidant and antimicrobial activities of the pulp and seed of *Anisophyllea laurina* R. Br. ex Sabine fruits. *Asian Pac. J. Trop. Biomed.*, **2016**, *6*, 20-25.  
<http://dx.doi.org/10.1016/j.apjtb.2015.09.023>
- [117] Gucwa, K.; Milewski, S.; Dymerski, T.; Szwed, P. Investigation of the Antifungal Activity and Mode of Action of *Thymus vulgaris*, *Citrus limonum*, *Pelargonium graveolens*, *Cinnamomum cassia*, *Ocimum basilicum*, and *Eugenia caryophyllus* Essential Oils. *Molecules*, **2018**, *23*(5), 1116.  
<http://dx.doi.org/10.3390/molecules23051116> PMID: 29738503
- [118] Baldim, I.; Tonani, L.; von Zeska Kress, M.R.; Pereira Oliveira, W. *Lippia sidoides* essential oil encapsulated in lipid nanosystem as an anti-*Candida* agent. *Ind. Crops Prod.*, **2019**, *127*, 73-81.  
<http://dx.doi.org/10.1016/j.indcrop.2018.10.064>
- [119] Almeida, E.T.D.C.; de Souza, G.T.; de Sousa Guedes, J.P.; Barbosa, I.M.; de Sousa, C.P.; Castellano, L.R.C.; Magnani, M.; de Souza, E.L. *Mentha piperita* L. essential oil inactivates spoilage yeasts in fruit juices through the perturbation of different physiological functions in yeast cells. *Food Microbiol.*, **2019**, *82*, 20-29.  
<http://dx.doi.org/10.1016/j.fm.2019.01.023> PMID: 31027774
- [120] Sadowska, B.; Budzyńska, A.; Stochmal, A.; Żuchowski, J.; Różalska, B. Novel properties of *Hippophae rhamnoides* L. twig and leaf extracts - anti-virulence action and synergy with antifungals studied *in vitro* on *Candida* spp. model. *Microb. Pathog.*, **2017**, *107*, 372-379.  
<http://dx.doi.org/10.1016/j.micpath.2017.04.020> PMID: 28428132
- [121] Lee, H.S.S.; Kim, Y. *Paeonia lactiflora* Inhibits Cell Wall Synthesis and Triggers Membrane Depolarization in *Candida albicans*. *J. Microbiol. Biotechnol.*, **2017**, *27*(2), 395-404.  
<http://dx.doi.org/10.4014/jmb.1611.11064> PMID: 28100900
- [122] Morais-Braga, M.F.; Carneiro, J.N.; Machado, A.J.; Sales, D.L.; Dos Santos, A.T.; Boligon, A.A.; Athayde, M.L.; Menezes, I.R.; Souza, D.S.; Costa, J.G.; Coutinho, H.D. Phenolic composition and medicinal usage of *Psidium guajava* Linn.: Antifungal activity or inhibition of virulence? *Saudi J. Biol. Sci.*, **2017**, *24*(2), 302-313.  
<http://dx.doi.org/10.1016/j.sjbs.2015.09.028> PMID: 28149166
- [123] Unuofin, J.O.; Otunola, G.A.; Afolayan, A.J. Phytochemical screening and *in vitro* evaluation of antioxidant and antimicrobial activities of *Kedrostis africana* (L.). *Cogn. Asian Pac. J. Trop. Biomed.*, **2017**, *7*, 901-908.  
<http://dx.doi.org/10.1016/j.apjtb.2017.09.008>
- [124] Bhuyan, D.J.; Vuong, Q.V.; Chalmers, A.C.; van Altena, I.A.; Bowyer, M.C.; Scarlett, C.J. Phytochemical, antibacterial and antifungal properties of an aqueous extract of *Eucalyptus microcorys* leaves. *S. Afr. J. Bot.*, **2017**, *112*, 180-185.  
<http://dx.doi.org/10.1016/j.sajb.2017.05.030>
- [125] Dal Piaz, F.; Bader, A.; Malafrente, N.; D'Ambola, M.; Petrone, A.M.; Porta, A.; Ben Hadda, T.; De Tommasi, N.; Bisio, A.; Severino, L. Phytochemistry of compounds isolated from the leaf-surface extract of *Psidium punctulata* (DC.) Vatke growing in Saudi Arabia. *Phytochemistry*, **2018**, *155*, 191-202.  
<http://dx.doi.org/10.1016/j.phytochem.2018.08.003> PMID: 30149245
- [126] Choi, J.H.; Kim, J.Y.; Jeong, E.T.; Choi, T.H.; Yoon, T.M. Preservative effect of *Camellia sinensis* (L.) Kuntze seed extract in soy sauce and its mutagenicity. *Food Res. Int.*, **2018**, *105*, 982-988.  
<http://dx.doi.org/10.1016/j.foodres.2017.11.059> PMID: 29433297
- [127] Morais-Braga, M.F.B.; Sales, D.L.; Carneiro, J.N.P.; Machado, A.J.T.; Dos Santos, A.T.L.; de Freitas, M.A.; Martins, G.M.A.B.; Leite, N.F.; de Matos, Y.M.L.S.; Tintino, S.R.; Souza, D.S.L.; Menezes, I.R.A.; Ribeiro-Filho, J.; Costa, J.G.M.; Coutinho, H.D.M. *Psidium guajava* L. and *Psidium brownianum* Mart ex DC.: Chemical composition and anti-*Candida* effect in association with fluconazole. *Microb. Pathog.*, **2016**, *95*, 200-207.  
<http://dx.doi.org/10.1016/j.micpath.2016.04.013> PMID: 27085299
- [128] Joshi, T.; Jain, T.; Mahar, R.; Singh, S.K.; Srivastava, P.; Shukla, S.K.; Mishra, D.K.; Bhatta, R.S.; Banerjee, D.; Kanjjiya, S. Pyranocarbazoles from *Murraya koenigii* (L.) Spreng. as antimicrobial agents. *Nat. Prod. Res.*, **2018**, *32*(4), 430-434.  
<http://dx.doi.org/10.1080/14786419.2017.1308363> PMID: 28368664
- [129] Rajkowska, K.; Kunicka-Styczyńska, A.; Maroszyńska, M. Selected Essential Oils as Antifungal Agents Against Antibiotic-Resistant *Candida* spp.: *In Vitro* Study on Clinical and Food-Borne Isolates. *Microb. Drug Resist.*, **2017**, *23*(1), 18-24.  
<http://dx.doi.org/10.1089/mdr.2016.0001> PMID: 27092733
- [130] Mtunzi, F.M.; Ejidike, I.P.; Ledwaba, I.; Ahmed, A.; Pakade, V.E.; Klink, M.J.; Modise, S.J.; Modise, S.J. Solvent-solvent fractionations of *Combretum erythrophyllum* (Burch.) leave extract: Studies of their antibacterial, antifungal, antioxidant and cytotoxicity potentials. *Asian Pac. J. Trop. Med.*, **2017**, *10*(7), 670-679.  
<http://dx.doi.org/10.1016/j.apjtm.2017.07.007> PMID: 28870343
- [131] Isa, A.I.; Awouafack, M.D.; Dzoyem, J.P.; Aliyu, M.; Magaji, R.A.; Ayo, J.O.; Eloff, J.N. Some *Strychnos spinosa* (Loganiaceae) leaf extracts and fractions have good antimicrobial activities and low cytotoxicities. *BMC Complement. Altern. Med.*, **2014**, *14*, 456.

- <http://dx.doi.org/10.1186/1472-6882-14-456> PMID: 25428165
- [132] Megeressa, M.; Bisrat, D.; Mazumder, A.; Asres, K. Structural elucidation of some antimicrobial constituents from the leaf latex of *Aloe trigonantha* L.C. Leach. *BMC Complement. Altern. Med.*, **2015**, *15*, 270.  
<http://dx.doi.org/10.1186/s12906-015-0803-4> PMID: 26264241
- [133] Rath, C.C.; Mohapatra, S. Susceptibility characterisation of *Candida* spp. to four essential oils. *Indian J. Med. Microbiol.*, **2015**, *33*(Suppl.), 93-96.  
<http://dx.doi.org/10.4103/0255-0857.150903> PMID: 25657164
- [134] Anibijuwon, I.; Gbala, I.; Abioye, J. Susceptibility of selected multi-drug resistant clinical isolates to leaves of *Carpolobia lutea*. *Ethiop. J. Health Sci.*, **2018**, *28*(2), 117-124.  
<http://dx.doi.org/10.4314/ejhs.v28i2.3> PMID: 29983509
- [135] Butassi, E.; Svetaz, L.A.; Ivancovich, J.J.; Feresin, G.E.; Tapia, A.; Zucchini, S.A. Synergistic mutual potentiation of antifungal activity of *Zuccagnia punctata* Cav. and *Larrea nitida* Cav. extracts in clinical isolates of *Candida albicans* and *Candida glabrata*. *Phyto-medicine*, **2015**, *22*(6), 666-678.  
<http://dx.doi.org/10.1016/j.phymed.2015.04.004> PMID: 26055132
- [136] Moreno, M.A.; Córdoba, S.; Zampini, I.C.; Mercado, M.I.; Pones-sa, G.; Alberto, M.R.; Nader-Macias, M.E.F.; Sayago, J.; Burgos-Edwards, A.; Schmeda-Hirschmann, G.; Isla, M.I. *Tetraglochin andina* Ciald.: A medicinal plant from the Argentinean highlands with potential use in vaginal candidiasis. *J. Ethnopharmacol.*, **2018**, *216*, 283-294.  
<http://dx.doi.org/10.1016/j.jep.2018.01.001> PMID: 29307753
- [137] Mahboubi, M.; Kazempour, N. The anti-candidal activity of *Satureja khuzistanica* ethanol extract against clinical isolates of *C. albicans*. *J. Mycol. Med.*, **2016**, *26*(1), e6-e10.  
<http://dx.doi.org/10.1016/j.mycmed.2015.11.003> PMID: 26849903
- [138] Boroja, T.; Mihailović, V.; Katanić, J.; Pan, S.P.; Nikles, S.; Im-bimbo, P.; Monti, D.M.; Stanković, N.; Stanković, M.S.; Bauer, R. The biological activities of roots and aerial parts of *Alchemilla vul-garis* L. *S. Afr. J. Bot.*, **2018**, *116*, 175-184.  
<http://dx.doi.org/10.1016/j.sajb.2018.03.007>
- [139] Zomorodian, K.; Saharkhiz, J.; Pakshir, K.; Immeripour, Z.; Sada-tsharif, A. The composition, antibiofilm and antimicrobial activi-ties of essential oil of *Ferula assa-foetida* oleo-gum-resin. *Biocatal. Agric. Biotechnol.*, **2018**, *14*, 300-304.  
<http://dx.doi.org/10.1016/j.bcab.2018.03.014>
- [140] Oliveira, J.R.; de Jesus Viegas, D.; Martins, A.P.R.; Carvalho, C.A.T.; Soares, C.P.; Camargo, S.E.A.; Jorge, A.O.C.; de Oliveira, L.D. *Thymus vulgaris* L. extract has antimicrobial and anti-inflammatory effects in the absence of cytotoxicity and genotoxicity. *Arch. Oral Biol.*, **2017**, *82*, 271-279.  
<http://dx.doi.org/10.1016/j.archoralbio.2017.06.031> PMID: 28683409
- [141] Sardi, J.C.; Freires, I.A.; Lazarini, J.G.; Infante, J.; de Alencar, S.M.; Rosalen, P.L. Unexplored endemic fruit species from Brazil: Antibiofilm properties, insights into mode of action, and systemic toxicity of four *Eugenia* spp. *Microb. Pathog.*, **2017**, *105*, 280-287.  
<http://dx.doi.org/10.1016/j.micpath.2017.02.044> PMID: 28259673
- [142] Forester, S.C.; Lambert, J.D. The role of antioxidant versus pro-oxidant effects of green tea polyphenols in cancer prevention. *Mol. Nutr. Food Res.*, **2011**, *55*(6), 844-854.  
<http://dx.doi.org/10.1002/mnfr.201000641> PMID: 21538850
- [143] de Castro, R.D.; de Souza, T.M.; Bezerra, L.M.; Ferreira, G.L.; Costa, E.M.; Cavalcanti, A.L. Antifungal activity and mode of ac-tion of thymol and its synergism with nystatin against *Candida* spe-cies involved with infections in the oral cavity: an *in vitro* study. *BMC Complement. Altern. Med.*, **2015**, *15*, 417.  
<http://dx.doi.org/10.1186/s12906-015-0947-2> PMID: 26601661
- [144] Chevalier, M.; Medioni, E.; Prêcheur, I. Inhibition of *Candida albicans* yeast-hyphal transition and biofilm formation by *Solidago virgaurea* water extracts. *J. Med. Microbiol.*, **2012**, *61*(Pt 7), 1016-1022.  
<http://dx.doi.org/10.1099/jmm.0.041699-0> PMID: 22422572
- [145] Filho, A.; Oliveira, H.; Sousa, J.; Meireles, D.R.P.; Maia, G.; Filho, J.M.B.; de Siquiera, J.P.J.; Lima, E. *In vitro* anti-*Candida* activity and mechanism of action of the flavonoid isolated from *Praxelis clematidea* against *Candida albicans* species. *J. Appl. Pharm. Sci.*, **2016**, *6*, 066-069.
- [146] Bernatoniene, J.; Kopustinskiene, D.M. The Role of Catechins in Cellular Responses to Oxidative Stress. *Molecules*, **2018**, *23*(4), 965.  
<http://dx.doi.org/10.3390/molecules23040965> PMID: 29677167
- [147] Rao, A.; Zhang, Y.; Muend, S.; Rao, R. Mechanism of antifungal activity of terpenoid phenols resembles calcium stress and inhibi-tion of the TOR pathway. *Antimicrob. Agents Chemother.*, **2010**, *54*(12), 5062-5069.  
<http://dx.doi.org/10.1128/AAC.01050-10> PMID: 20921304
- [148] Karygianni, L.; Al-Ahmad, A.; Argyropoulou, A.; Hellwig, E.; Anderson, A.C.; Skaltsounis, A.L. Natural antimicrobials and oral microorganisms: A systematic review on herbal interventions for the eradication of multispecies oral biofilms. *Front. Microbiol.*, **2016**, *6*, 1529.  
<http://dx.doi.org/10.3389/fmicb.2015.01529> PMID: 26834707
- [149] The Centre for Evidence-Based Medicine. <https://www.cebm.net/2020>.
- [150] Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D.G. PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.*, **2009**, *6*(7)e1000097  
<http://dx.doi.org/10.1371/journal.pmed.1000097> PMID: 19621072
- [151] Azwanida, N.N. A Review on the Extraction Methods Use in Medicinal Plants, Principle, Strength and Limitation. *Med. Aromat. Plants*, **2015**, *4*, 3.