

# Allergic Rhinitis and its Impact on Asthma Update (ARIA 2008) The Perspective From Spain

J Mullol,<sup>1,2,8</sup> A Valero,<sup>2,3,8</sup> I Alobid,<sup>1,2,8</sup> J Bartra,<sup>2,3,8</sup> AM Navarro,<sup>4</sup> T Chivato,<sup>5</sup>  
N Khaltaev,<sup>6</sup> J Bousquet<sup>7,8</sup>

<sup>1</sup> Rhinology Unit & Smell Clinic, ENT Department, Hospital Clínic – IDIBAPS, Barcelona, Catalonia, Spain

<sup>2</sup> Centre for Biomedical Investigation in Respiratory Diseases (CIBERES), Spain

<sup>3</sup> Allergy Unit, Pneumology and Allergy Department, Hospital Clínic – IDIBAPS, Barcelona, Catalonia, Spain

<sup>4</sup> Allergy Unit, Hospital El Tomillar, AH de Valme, Seville, Spain

<sup>5</sup> Immunology and Allergy Department, Hospital of the Air, Madrid, Spain

<sup>6</sup> GARD/ARIA Coordinator, Geneva, Switzerland

<sup>7</sup> University Hospital and INSERM, Hôpital Arnaud de Villeneuve, Montpellier, France

<sup>8</sup> Global Allergy and Asthma European Network (GA2LEN)

## ■ Abstract

Allergic rhinitis is a global health problem. Over 600 million patients suffer from this disease worldwide. ARIA (Allergic Rhinitis and its Impact on Asthma), an evidence-based document, was produced and published in 2001 using an extensive review of the available literature. The ARIA 2008 update was recently published and covers tertiary prevention of allergy, pharmacologic treatments, and immunotherapy. Nonallergic rhinitis is still a controversial area and may pose some treatment problems. Another important aspect of the ARIA update is the comorbidity of allergic rhinitis, in particular, asthma. The recommendations of the 2008 ARIA Update, as in 2001, are that patients with allergic rhinitis, particularly if persistent, should be evaluated for asthma, patients with asthma should be evaluated for rhinitis, and an effective and safe combination strategy should be used to treat diseases of the upper and lower airways. Over the last few years, several studies performed in Spain report new data on the prevalence of allergic rhinitis, sensitivity to common aeroallergens, comorbidity of allergic rhinitis and asthma, and impact on quality of life. The studies reviewed in this manuscript confirm—as do those from other developed countries—the enormous impact of the disease on society and health care in Spain.

**Key words:** Allergic rhinitis. Epidemiology. Comorbidities. Quality of life.

## ■ Resumen

La rinitis alérgica constituye un problema de salud global. Más de 600 millones de pacientes sufren esta enfermedad en todo el mundo. ARIA (Allergic Rhinitis and its Impact on Asthma), un documento basado en pruebas científicas, fue elaborado y publicado en 2001 mediante una revisión sistemática de la literatura disponible. La actualización 2008 de ARIA ha sido publicada recientemente y una amplia lista de terapias han sido abordadas y actualizadas en cuanto a la prevención de la alergia, el tratamiento farmacológico y la inmunoterapia. La rinitis no alérgica constituye todavía un tema de discusión pudiendo ofrecer problemas de tratamiento. Otro aspecto importante de la actualización de ARIA son las comorbilidades de la rinitis alérgica, el asma en particular. Las recomendaciones de la Actualización 2008 de ARIA, como en 2001, son que en los pacientes con rinitis alérgica, sobre todo si es persistente, debe estudiarse la presencia de asma; en los pacientes con asma debe investigarse la rinitis; y debe emplearse una estrategia para el tratamiento combinado de la rinitis y el asma en términos de eficacia y gravedad. Durante los últimos años se han realizado en España diversos estudios que han aportado nuevos datos sobre la prevalencia de la rinitis alérgica en nuestra población, la sensibilidad a aeroalérgenos, la comorbilidad de rinitis alérgica y asma, y el impacto sobre la calidad de vida. Estos estudios, revisados en este artículo, confirman, al igual que en otros países desarrollados, el enorme impacto social y sanitario que la rinitis tiene en España.

**Palabras clave:** Rinitis alérgica. Epidemiología. Comorbilidades. Calidad de vida.

## ARIA Update (2008)

Allergic rhinitis is a symptomatic disorder of the mucous membrane of the nose induced after allergen exposure due to an IgE-mediated inflammation of the membranes lining the nose. It was defined in 1929: "The three cardinal symptoms in nasal reactions occurring in allergy are sneezing, nasal obstruction and mucous discharge" [1].

Allergic rhinitis is a global health problem that affects patients of all ages and ethnic groups. It causes major illness and disability worldwide. Allergic rhinitis affects social life, sleep, and performance at school and work [2-5], and its economic impact is substantial. However, rhinitis is still underdiagnosed and undertreated [6].

Over 600 million patients suffer from this disease [7-10], but there are still differences between rural and urban areas, both in developed and developing countries [8,10-12], possibly due to differences in immune reactions [14].

In 1999, during the ARIA (Allergic Rhinitis and its Impact on Asthma) World Health Organization (WHO) workshop, an evidence-based document was produced using an extensive review of the literature available up to December 1999 [15]. The statements of evidence for the development of ARIA followed WHO rules and the recommendations of Shekelle et al [16].

The ARIA document presented the state-of-the-art for the specialist, general practitioner, and other health care professionals. Its aims were as follows:

- To update knowledge of allergic rhinitis
- To highlight the impact of allergic rhinitis on asthma
- To provide an evidence-based documented review of diagnostic methods
- To provide an evidence-based review of the treatments available
- To propose a stepwise approach to the management of the disease

However, an update of the ARIA guidelines was necessary for the following reasons:

- The large number of papers published over the last 7 years has increased our knowledge [17-22].
- The ARIA classification was proposed by an expert group and needed to be validated in terms of classification and management [14]. New studies showed consistently that "intermittent" and "persistent" are not synonymous with "seasonal" and "perennial" [23,24]. Several reports have now validated this classification [25,26], although some authors proposed extending the severity of allergic rhinitis to 3 levels [27,28]. However, since this would not lead to a difference in treatment, the ARIA experts proposed continuing to classify the severity of rhinitis as "mild" and "moderate/severe."
- New methods of diagnosis have been proposed for allergic and nonallergic rhinitis [29-32]. The diagnosis of allergic rhinitis is often easy, but in some cases it is problematic and many patients are still underdiagnosed, often because they do not perceive the symptoms of rhinitis as a disease.
- Gaps in our knowledge in the first ARIA document have now been filled. These include complementary and alternative medicine [18], sports and rhinitis in athletes [19,33,34], and rhinitis and its links with asthma in children [35-39].

The ARIA update began in 2004 and was published earlier this year [42]. Several chapters were extensively reviewed using the Shekelle evidence-based model [16], and papers published in peer-reviewed journals [17-22]. These papers cover the area of tertiary prevention of allergy, complementary and alternative medicine, pharmacotherapy and anti-IgE treatment, allergen-specific immunotherapy, links between rhinitis and asthma, and mechanisms of rhinitis. The need arose for a global document to highlight the interactions between the upper and the lower airways including diagnosis, epidemiology, common risk factors, management, and prevention. Moreover, attention was also given to allergy in developing countries [40,41].

The grading of evidence and the recommendation for an evidence-based management system in the ARIA 2008 update did not follow the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach [43, 44]. It is expected that some of the recommendations offered by the 2008 ARIA update would differ if the GRADE approach had been applied.

A large list of treatments was considered in the ARIA 2008 update [42]. Intranasal corticosteroids are the first-line therapy in patients with moderate to severe disease and are also effective against ocular symptoms [45]. H<sub>1</sub>-antihistamines are important treatments for all patients and leukotriene receptor antagonists are particularly important for patients with rhinitis and asthma [46,47]. Tertiary prevention of allergy is still a matter of debate, since clinical trials do not usually show any efficacy of single allergen avoidance measures [17]. Sublingual immunotherapy has proven to be a safe and effective treatment [48-51], but clinical trials need to be standardized [52,53]. An algorithm of the management of allergic rhinitis is provided (Figure). However, our progress in understanding the mechanisms of allergic rhinitis is continuous and novel treatment approaches are constantly being published [54].

Nonallergic rhinitis is still a matter of discussion [55] and may pose some treatment problems [56].

Another important aspect of ARIA was to consider comorbid conditions of allergic rhinitis, in particular, asthma. Epidemiologic studies throughout the world have consistently shown that asthma and rhinitis often coexist in the same patient [57-60]. The vast majority of patients with asthma have rhinitis, but the prevalence of asthma in rhinitis patients still needs to be assessed [61,62]. The treatment of nasal symptoms has little effect on the lower airways, but there have been some compelling data suggesting that new studies with innovative methods need to be started [63,64]. Specific immunotherapy in patients with allergic rhinitis has a prolonged effect on the development of asthma when stopped [65].

The perception of patients and physicians regarding the links between asthma and rhinitis varies between countries, but this perception appears to be stronger than expected [66,67]. However, knowledge is not directly translated into practice, since fewer physicians coprescribe treatments for rhinitis and asthma.

The recommendations of the ARIA workshop in 1999 are still valid [15], and patients with allergic rhinitis, in particular persistent allergic rhinitis, should be evaluated for asthma. Patients with asthma should be evaluated for rhinitis, and an efficacious and combined strategy should be adopted to treat diseases of the upper and lower airway.




Figure. Management (diagnosis and treatment) scheme of allergic rhinitis according to the 2008 ARIA Update (with permission from reference 42). Abbreviations: CS, glucocorticosteroid; LTRA, leukotriene receptor antagonist.

## Allergic Rhinitis: the Perspective From Spain

Over the last few years, several Spanish studies have reported new data on the prevalence of allergic rhinitis, sensitivity to common aeroallergens, comorbidity of allergic rhinitis and asthma, and impact on quality of life. The results of these studies confirm (as is the case in other developed countries) the enormous impact of the disease on society and health care.

Bauchau and Durham [23] performed a pan-European epidemiological study on the prevalence of allergic rhinitis in 6 European countries (Belgium, France, Germany, Italy,

Spain, and the United Kingdom). The study was performed in 2 phases: during the first phase, 9646 subjects were surveyed by phone and, during the second, 725 subjects were selected and included in a clinical trial in which 411 subjects were diagnosed with allergic rhinitis.

The Estudio Ibérico [68], was a descriptive, observational, cross-sectional, population-based study carried out in Portugal and Spain in a population of allergic rhinitis patients to evaluate skin sensitization, characteristics of rhinitis, and comorbidity of rhinitis and asthma. The sample was composed of patients aged 10 to 50 years with allergic rhinitis consecutively attended at allergology centers. Patients were diagnosed using skin prick test to 20 standardized aeroallergens, and rhinitis was classified

according to ARIA and the etiological agent and asthma according to the Global Initiative for Asthma (GINA) [69].

The Spanish Society of Allergology and Clinical Immunology (SEAIC) developed 2 epidemiological studies, Alergológica, with the objective of obtaining information on allergic patients attended at outpatient allergology clinics. Those surveys were performed in 1992 and again in 2005 with the inclusion of 4005 and 4991 patients, respectively [70,71].

Two other prospective studies [72,73] have recently been performed in allergology (Oneair) and pneumology (Rinair) outpatient clinics to study the prevalence of allergic rhinitis among patients with diagnosed asthma from 2005 to 2006. Using an adequate geographic distribution, 170 specialists in allergology and 172 in pneumology participated in the studies. Demographic data, outcome of asthma and rhinitis, and classification according to GINA and ARIA were collected from 942 (Oneair) and 703 (Rinair) asthma patients.

To develop and validate a specific Spanish questionnaire for allergic rhinitis, the ESPRINT (ESTudio de la calidad de vida en Pacientes con RINOconjuntivitis) study was performed with the participation of over 40 centers and 800 patients, 413 in the item-reduction phase and 400 in the questionnaire-validation phase [74,75]. The inclusion criterion for patients with allergic rhinitis participating in the study was a total symptom score greater than 3 (score of 0 to 3 in each of nasal congestion, rhinorrhea, nasal itching, and sneezing). In the context of this project, new criteria to classify the severity of allergic rhinitis as mild, moderate, and severe according to ARIA severity items were also established [28].

### Epidemiology

The pan-European study reported a global prevalence of allergic rhinitis of 22.7% (95% confidence interval [95% CI]; 21.1-22.2) with a variation in European countries ranging from 16.9% (95% CI, 12.9-20.9) for Italy to 28.5% (95% CI, 24.5-32.5) for Belgium [23]. In the global survey, 45% of cases had no previous diagnosis (underdiagnosis) of allergic rhinitis. Five centers and cities (Barcelona, Bilbao, Madrid, Sevilla, Valencia) participated for Spain. In the first phase, 1600 telephone surveys were performed and, in the second phase, 100 subjects were studied to confirm the diagnosis of allergic rhinitis. The prevalence of allergic rhinitis in Spain was 21.5% (18.5-24.4).

The Estudio Ibérico studied 3225 subjects (2020 for Spain and 1205 for Portugal) [67] with a mean age of 27 years. Major house dust mites (*Dermatophagoides pteronyssinus*, *Dermatophagoides farinae*) and grass and olive pollens were the most prevalent allergens, with skin polysensitization accounting for 95% of cases with a mean (SD) of 6.5 (3.7) sensitizations per subject. There were more cases of sensitization to pollens (n=2298) and house dust mites (n=2245) than to animal dander (n=1329) and mold (*Alternaria*) (n=399). Monosensitization accounted for 26% of cases for pollens and house dust mites and 2% of cases for animal dander and molds, while allergen group monosensitization accounted for 37% of cases (18% for pollens, 18% for house dust mites, 0.8% for animal dander, and 0.3% for *Alternaria*).

In the 1992 and 2005 Alergológica surveys [70,71], the prevalence of diagnosed allergic rhinitis was 52% and 55%, respectively, and 47% for the pediatric population (1992). The mean (SD) age of rhinoconjunctivitis patients was 32.1 (18) and 29.9 (15), respectively.

### Classification of Allergic Rhinitis According to the ARIA Criteria

Several studies performed in Spain over recent years have investigated the new classification of allergic rhinitis according to the ARIA criteria [15,23-26]. In the pan-European study, 21% (13.5-30.3) of allergic patients were classified as persistent in Spain while 29.3% (24.9-34.0) were persistent in the global study [23]. In the Estudio Ibérico, 64% of subjects with allergic rhinitis who visited allergy outpatient clinics (N=3225) were classified as persistent and 41% as moderate-to-severe [68].

In the Oneair study, 54% of asthmatic patients with allergic rhinitis were classified as persistent and 57% as moderate-to-severe (24% mild intermittent, 22% moderate/severe intermittent, 19% mild persistent, and 35% moderate/severe persistent) [72]. In the Rinair study, 43% of patients were classified as persistent and 74% as moderate-to-severe (9.2% mild intermittent, 47.4% moderate/severe intermittent, 16.3% mild persistent, and 27.1% moderate/severe persistent) [73].

In the ESPRINT study, 59% of the patients consulted with allergic rhinitis (N=400) in a specialist (allergology, otorhinolaryngology) outpatient clinic were classified as persistent [74,75]. To differentiate between moderate and severe allergic rhinitis, a new criterion was reported according to ARIA severity items [28]: patients with 1, 2, or 3 severity items affected continue to be classified as moderate, whereas patients with all 4 items affected are now classified as severe.

### Comorbidity of Rhinitis and Asthma

Most studies on the association between rhinitis and asthma evaluate the prevalence of asthma in patients with allergic rhinitis, while few assess the prevalence of rhinitis in patients with asthma. The prevalence of allergic rhinitis among asthma patients varies widely in published studies, ranging from 80% to 95% in the most recent [76]. In the Estudio Ibérico, 49% of patients with allergic rhinitis had concomitant asthma (56% intermittent, 44% persistent, 33% mild, 10% moderate, and 1% severe [68]). In the 2005 Alergológica study, 37% of patients with allergic rhinitis had concomitant asthma and 65% conjunctivitis [69].

In the Oneair (allergologists) and Rinair (pneumologists) studies, the prevalence of allergic rhinitis among asthma patients was 71% and 89.5%, respectively [72,73]. However, it is interesting to emphasize that, in the Rinair study, the prevalence was much higher among allergic (84%) than nonallergic (51%) asthmatics. In the Oneair study, the prevalence of asthma severity groups according to the GINA classification was 38.5% intermittent and 61.5% persistent (29.4% mild, 27.2% moderate, and 4.9% severe). In the Rinair study, the prevalence of asthma by severity was 24.5% intermittent and 75.5% persistent (35.4% mild, 32.7% moderate, and 7.4% severe).

In both studies, there was a significant correlation ( $P < .0001$ ) between the severity of rhinitis and of asthma. The prevalence of rhinitis decreased among older and more severely affected asthmatic patients.

Of the 400 allergic rhinitis patients for whom the ESPRINT questionnaire was validated, 39% had concomitant asthma and 53% conjunctivitis [74,75].

### Quality of Life

During the last few decades, considerable importance has been given not only to the symptoms present in a variety of diseases, but also to how these diseases affect the quality of life (QoL) of patients. Thus, generic and specific questionnaires have been developed to assess the QoL of allergic rhinitis patients [77-81].

The 2005 Alergológica study, which used the SF-12 generic questionnaire that assesses the impact on health-related QoL [82], showed that values for allergic rhinitis patients were located between the 25th percentile (physical component) and 20th percentile (mental component) for the general population of Spain, meaning that the QoL perceived by patients with allergic rhinitis was lower than that perceived by 75% of the general population [71].

In the ESPRINT study, a new specific questionnaire to assess the QoL in allergic rhinitis patients was developed and validated for the Spanish population. The ESPRINT questionnaire has 2 versions, a long version with 28 items to be used in research studies, and a short version with 15 items to be used in daily clinical practice. Using both long and short questionnaires has revealed an impact of allergic rhinitis on QoL, and this impact is higher in persistent than in intermittent allergic rhinitis. There was a significant correlation with the severity of allergic rhinitis according to the ARIA classification, which made it possible to differentiate between different levels of rhinitis severity [28,74,75].

### Socioeconomic Burden

Few studies have evaluated the economic and social burden of allergic rhinitis [2, 4, 83-85]. In Spain, no studies have been performed to date on the economic burden of allergic rhinitis. In the 2005 Alergológica study [71], 61% of patients visited their family physician during the last quarter with an average of 2.1 visits. During the last year, 29% of patients visited a specialist with an average of 1.7 visits, while 22% visited the emergency room, with an average of 1.9 visits per year (1.2% of patients had to be hospitalized). In allergic rhinitis/conjunctivitis patients, 6% reported absenteeism from work related to their disease, 66% with only 1 episode, and an average duration of 15.6 days. Among a population sample of 337 students, an average of 8 working days lost during the last year was reported. Work absenteeism by parents was reported in 15% of a pediatric sample (N=513), with an average of 4 days/year. In general, good school performance was reported for 79% of children.

In conclusion, due to the high prevalence of allergic rhinitis, its impact on quality of life, and the social and health care burden, further investigation is necessary to obtain pharmaco-economic data on allergic rhinitis in Spain.

## References

- Hansel F. Clinical and histopathologic studies of the nose and sinuses in allergy. *J Allergy*. 1929;1:43-70.
- Canonica GW, Bousquet J, Mullol J, Scadding GK, Virchow JC. A survey of the burden of allergic rhinitis in Europe. *Allergy*. 2007;62 (Suppl 85):17-25.
- van Oene CM, van Reijl EJ, Sprangers MA, Fokkens WJ. Quality-assessment of disease-specific quality of life questionnaires for rhinitis and rhinosinusitis: a systematic review. *Allergy*. 2007;62(12):1359-71.
- Schatz M. A survey of the burden of allergic rhinitis in the USA. *Allergy*. 2007;62(Suppl 85):9-16.
- Walker S, Khan-Wasti S, Fletcher M, Cullinan P, Harris J, Sheikh A. Seasonal allergic rhinitis is associated with a detrimental effect on examination performance in United Kingdom teenagers: case-control study. *J Allergy Clin Immunol*. 2007;120(2):381-7.
- Maurer M, Zuberbier T. Undertreatment of rhinitis symptoms in Europe: findings from a cross-sectional questionnaire survey. *Allergy* 2007;62(9):1057-63.
- Bousquet J, Dahl R, Khaltaev N. Global alliance against chronic respiratory diseases. *Allergy*. 2007;62(3):216-23.
- Bousquet J, Khaltaev N. Global surveillance, prevention and control of chronic respiratory diseases. A comprehensive approach. *Global Alliance Against Chronic Respiratory Diseases*. World Health Organization. ISBN 978 92 4 156346 8. 2007:148 pages.
- Asher MI, Montefort S, Bjorksten B, Lai CK, Strachan DP, Weiland SK, Williams H; ISAAC Phase Three Study Group. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet*. 2006;368(9537):733-43.
- Ait-Khaled N, Odhiambo J, Pearce N, Adjoh KS, Maesano IA, Benhabyles B, Bouhayad Z, Bahati E, Camara L, Catteau C, El Sony A, Esamai FO, Hypolite IE, Melaku K, Musa OA, Ng'ang'a L, Onadeko BO, Saad O, Jerray M, Kayembe JM, Koffi NB, Khaldi F, Kuaban C, Voyi K, M'Boussa J, Sow O, Tidjani O, Zar HJ. Prevalence of symptoms of asthma, rhinitis and eczema in 13- to 14-year-old children in Africa: the International Study of Asthma and Allergies in Childhood Phase III. *Allergy*. 2007;62(3):247-58.
- Pekkarinen PT, von Hertzen L, Laatikainen T, Makela MJ, Jousilahti P, Kosunen TU, Pantelejev V, Vartiainen E, Haahela T. A disparity in the association of asthma, rhinitis, and eczema with allergen-specific IgE between Finnish and Russian Karelia. *Allergy*. 2007;62(3):281-7.
- Majkowska-Wojciechowska B, Pelka J, Korzon L, Kozłowska A, Kaczala M, Jarzebska M, Gwardys T, Kowalski ML. Prevalence of allergy, patterns of allergic sensitization and allergy risk factors in rural and urban children. *Allergy*. 2007;62(9):1044-50.
- Viinanen A, Munhbayarlah S, Zevgee T, Narantsetseg L, Naidansuren T, Koskenvuo M, Helenius H, Terho EO. The protective effect of rural living against atopy in Mongolia. *Allergy*. 2007;62(3):272-80.
- van Ree R, Yazdanbakhsh M. Allergic disorders in African countries: linking immunology to accurate phenotype. *Allergy*. 2007;62(3):237-46.
- Bousquet J, Van Cauwenberge P, Khaltaev N. Allergic rhinitis

- and its impact on asthma. *J Allergy Clin Immunol.* 2001;108(5 Suppl):S147-334.
16. Shekelle PG, Woolf SH, Eccles M, Grimshaw J. Clinical guidelines: developing guidelines. *BMJ.* 1999;318(7183):593-6.
  17. Custovic A, Wijk RG. The effectiveness of measures to change the indoor environment in the treatment of allergic rhinitis and asthma: ARIA update (in collaboration with GA(2)LEN). *Allergy.* 2005;60(9):1112-5.
  18. Passalacqua G, Bousquet PJ, Carlsen KH, Kemp J, Lockey RF, Niggemann B, Pawankar R, Price D, Bousquet J. ARIA update: I--Systematic review of complementary and alternative medicine for rhinitis and asthma. *J Allergy Clin Immunol.* 2006;117(5):1054-62.
  19. Bonini S, Bonini M, Bousquet J, Brusasco V, Canonica GW, Carlsen KH, Corbetta L, Cummiskey J, Delgado L, Del Giacco SR, Haahtela T, Jaeger S, Moretti C, Palange P, Passalacqua G, Passali D, Pedersen BK, Popov T, Rasi G, Ventura MT, Vignola AM. Rhinitis and asthma in athletes: an ARIA document in collaboration with GA2LEN. *Allergy.* 2006;61(6):681-92.
  20. Bousquet J, van Cauwenberge P, Ait Khaled N, Bachert C, Baena-Cagnani CE, Bouchard J, Bunnag C, Canonica GW, Carlsen KH, Chen YZ, Cruz AA, Custovic A, Demoly P, Dubakiene R, Durham S, Fokkens W, Howarth P, Kemp J, Kowalski ML, Kvedariene V, Lipworth B, Lockey R, Lund V, Mavale-Manuel S, Meltzer EO, Mullol J, Naclerio R, Nekam K, Ohta K, Papadopoulos N, Passalacqua G, Pawankar R, Popov T, Potter P, Price D, Scadding G, Simons FE, Spicak V, Valovirta E, Wang DY, Yawn B, Yusuf O; GA2LEN. Pharmacologic and anti-IgE treatment of allergic rhinitis ARIA update (in collaboration with GALEN). *Allergy.* 2006;61(9):1086-96.
  21. Passalacqua G, Durham SR. Allergic rhinitis and its impact on asthma update: allergen immunotherapy. *J Allergy Clin Immunol.* 2007;119(4):881-91.
  22. Cruz AA, Popov T, Pawankar R, Annesi-Maesano I, Fokkens W, Kemp J, Ohta K, Price D, Bousquet J; ARIA Initiative Scientific Committee. Common characteristics of upper and lower airways in rhinitis and asthma: ARIA update, in collaboration with GA(2)LEN. *Allergy.* 2007;62 (Suppl 84):1-41.
  23. Bauchau V, Durham SR. Prevalence and rate of diagnosis of allergic rhinitis in Europe. *Eur Respir J.* 2004;24(5):758-64.
  24. Demoly P, Allaert FA, Lecasble M, Bousquet J. Validation of the classification of ARIA (allergic rhinitis and its impact on asthma). *Allergy.* 2003;58(7):672-5.
  25. Bachert C, van Cauwenberge P, Olbrecht J, van Schoor J. Prevalence, classification and perception of allergic and nonallergic rhinitis in Belgium. *Allergy.* 2006;61(6):693-8.
  26. Todo-Bom A, Loureiro C, Almeida MM, Nunes C, Delgado L, Castel-Branco G, Bousquet J. Epidemiology of rhinitis in Portugal: evaluation of the intermittent and the persistent types. *Allergy.* 2007;62(9):1038-43.
  27. Van Hoecke H, Vastesaeger N, Dewulf L, De Bacquer D, van Cauwenberge P. Is the allergic rhinitis and its impact on asthma classification useful in daily primary care practice? *J Allergy Clin Immunol.* 2006;118(3):758-9.
  28. Valero A, Ferrer M, Sastre J, Navarro AM, Monclus L, Marti-Guadano E, Herdman M, Dávila I, Del Cuvillo A, Colás C, Baró E, Antépara I, Alonso J, Mullol J. A new criterion by which to discriminate between patients with moderate allergic rhinitis and patients with severe allergic rhinitis based on the Allergic Rhinitis and its Impact on Asthma severity items. *J Allergy Clin Immunol.* 2007;120(2):359-65.
  29. Bousquet PJ, Combescure C, Neukirch F, Klossek JM, Mechin H, Daures JP, Bousquet J. Visual analog scales can assess the severity of rhinitis graded according to ARIA guidelines. *Allergy.* 2007;62(4):367-72.
  30. Boot JD, de Kam ML, Mascelli MA, Miller B, van Wijk RG, de Groot H, Cohen AF, Diamant Z. Nasal nitric oxide: longitudinal reproducibility and the effects of a nasal allergen challenge in patients with allergic rhinitis. *Allergy.* 2007;62(4):378-84.
  31. Juniper EF, Riis B, Juniper BA. Development and validation of an electronic version of the Rhinoconjunctivitis Quality of Life Questionnaire. *Allergy.* 2007;62(9):1091-3.
  32. Nizankowska-Mogilnicka E, Bochenek G, Mastalerz L, Swierczy ska M, Picado C, Scadding G, Kowalski ML, Setkowicz M, Ring J, Brockow K, Bachert C, Wöhrl S, Dahlén B, Szczeklik A. Aspirin provocation tests for diagnosis of aspirin sensitivity. EAACI/GA2LEN guideline. *Allergy.* 2007;62(10):1111-8.
  33. Bonini M, Lapucci G, Petrelli G, Todaro A, Pamich T, Rasi G, Bonini S. Predictive value of allergy and pulmonary function tests for the diagnosis of asthma in elite athletes. *Allergy.* 2007;62(10):1166-70.
  34. Bonini S, Rasi G, Brusasco V, Carlsen KH, Crimi E, Popov T, Schultze-Werninghaus G, Gramiccioni C, Bonini M, Passali D, Bachert C, van Cauwenberge PB, Bresciani M, Bonini S, Calonge M, Montan PG, Serapiao Dos Santos M, Belfort R Jr, Lambiase A, Sacchetti M. Nonspecific provocation of target organs in allergic diseases: EAACI-GA(2)LEN consensus report. *Allergy.* 2007;62(6):683-94.
  35. Burgess JA, Walters EH, Byrnes GB, Matheson MC, Jenkins MA, Wharton CL, Johns DP, Abramson MJ, Hopper JL, Dharmage SC. Childhood allergic rhinitis predicts asthma incidence and persistence to middle age: a longitudinal study. *J Allergy Clin Immunol* 2007;120(4):863-9.
  36. Chatkin MN, Menezes AM, Victora CG, Barros FC. High prevalence of asthma in preschool children in Southern Brazil: a population-based study. *Pediatric Pulmonol.* 2003;35(4):296-301.
  37. Giovannini M, Agostoni C, Riva E, Salvini F, Ruscitto A, Zuccotti GV, Giovannini M, Agostoni C, Riva E, Salvini F, Ruscitto A, Zuccotti GV, Radaelli G; Felicita Study Group. A randomized prospective double blind controlled trial on effects of long-term consumption of fermented milk containing *Lactobacillus casei* in pre-school children with allergic asthma and/or rhinitis. *Pediatr Res.* 2007;62(2):215-20.
  38. Viegli G, La Grutta S. Rhinoconjunctivitis and wheeze in preschool children: a different relationship than in adults (united or coexistent airways disease)? *Allergy.* 2007;62(4):344-7.
  39. Choi SH, Yoo Y, Yu J, Rhee CS, Min YG, Koh YY. Bronchial hyperresponsiveness in young children with allergic rhinitis and its risk factors. *Allergy.* 2007;62(9):1051-6.
  40. Bateman ED, Jithoo A. Asthma and allergy - a global perspective. *Allergy.* 2007;62(3):213-5.
  41. English RG, Fairall LR, Bateman ED. Keeping allergy on the agenda: integrated guidelines for respiratory disease in developing countries. *Allergy.* 2007;62(3):224-9.
  42. Bousquet J, Khaltaev N, Cruz AA, Denburg J, Fokkens WJ, Togias A, Zuberbier T, Baena-Cagnani CE, Canonica GW, van Weel C, Agache I, Ait-Khaled N, Bachert C, Blaiss MS, Bonini S, Boulet LP, Bousquet PJ, Camargos P, Carlsen KH, Chen Y, Custovic A,

- Dahl R, Demoly P, Douagui H, Durham SR, van Wijk RG, Kalayci O, Kaliner MA, Kim YY, Kowalski ML, Kuna P, Le LT, Lemiere C, Li J, Lockey RF, Mavale-Manuel S, Meltzer EO, Mohammad Y, Mullol J, Naclerio R, O'Hehir RE, Ohta K, Ouedraogo S, Palkonen S, Papadopoulos N, Passalacqua G, Pawankar R, Popov TA, Rabe KF, Rosado-Pinto J, Scadding GK, Simons FE, Toskala E, Valovirta E, van Cauwenberge P, Wang DY, Wickman M, Yawn BP, Yorgancioglu A, Yusuf OM, Zar H, Annesi-Maesano I, Bateman ED, Ben Kheder A, Boakye DA, Bouchard J, Burney P, Busse WW, Chan-Yeung M, Chavannes NH, Chuchalin A, Dolen WK, Emuzyte R, Grouse L, Humbert M, Jackson C, Johnston SL, Keith PK, Kemp JP, Klossek JM, Larenas-Linnemann D, Lipworth B, Malo JL, Marshall GD, Naspitz C, Nekam K, Niggemann B, Nizankowska-Mogilnicka E, Okamoto Y, Orru MP, Potter P, Price D, Stoloff SW, Vandenplas O, Viegi G, Williams D. ARIA Update. *Allergy*. 2008;63(Suppl 86):8-160.
43. Schunemann HJ, Hill SR, Kakad M, Vist GE, Bellamy R, Stockman L, Wisloff TF, Del Mar C, Hayden F, Uyeki TM, Farrar J, Yazdanpanah Y, Zucker H, Beigel J, Chotpitayasunondh T, Hien TT, Ozbay B, Sugaya N, Oxman AD. Transparent development of the WHO rapid advice guidelines. *PLoS Med*. 2007 May;4(5):e119.
  44. Brozek JL, Baena-Cagnani CE, Bonini S, Canonica GW, Rasi G, van Wijk RG, Zuberbier T, Guyatt G, Bousquet J, Schunemann HJ. Methodology for development of the Allergic Rhinitis and its Impact on Asthma guideline 2008 update. *Allergy*. 2008;63(1):38-46.
  45. Fokkens WJ, Jogi R, Reinartz S, Sidorenko I, Sitkauskienė B, van Oene C, Faris MA, Ellsworth A, Caldwell MF. Once daily fluticasone furoate nasal spray is effective in seasonal allergic rhinitis caused by grass pollen. *Allergy*. 2007;62(9):1078-84.
  46. Philip G, Nayak AS, Berger WE, Leynadier F, Vrijens F, Dass SB, Reiss TF. The effect of montelukast on rhinitis symptoms in patients with asthma and seasonal allergic rhinitis. *Curr Med Res Opin*. 2004;20(10):1549-58.
  47. Barnes ML, Menzies D, Fardon TC, Burns P, Wilson AM, Lipworth BJ. Combined mediator blockade or topical steroid for treating the unified allergic airway. *Allergy*. 2007;62(1):73-80.
  48. Durham SR, Riis B. Grass allergen tablet immunotherapy relieves individual seasonal eye and nasal symptoms, including nasal blockage. *Allergy*. 2007;62(8):954-7.
  49. Durham SR, Yang WH, Pedersen MR, Johansen N, Rak S. Sublingual immunotherapy with once-daily grass allergen tablets: a randomized controlled trial in seasonal allergic rhinoconjunctivitis. *J Allergy Clin Immunol*. 2006;117(4):802-9.
  50. Dahl R, Kapp A, Colombo G, de Monchy JG, Rak S, Emminger W, Rivas MF, Ribel M, Durham SR. Efficacy and safety of sublingual immunotherapy with grass allergen tablets for seasonal allergic rhinoconjunctivitis. *J Allergy Clin Immunol*. 2006;118(2):434-40.
  51. Didier A, Malling HJ, Worm M, Horak F, Jager S, Montagut A, André C, de Beaumont O, Melac M. Optimal dose, efficacy, and safety of once-daily sublingual immunotherapy with a 5-grass pollen tablet for seasonal allergic rhinitis. *J Allergy Clin Immunol*. 2007;120(6):1338-45.
  52. Canonica GW, Baena-Cagnani CE, Bousquet J, Bousquet PJ, Lockey RF, Malling HJ, Passalacqua G, Potter P, Valovirta E. Recommendations for standardization of clinical trials with Allergen Specific Immunotherapy for respiratory allergy. A statement of a World Allergy Organization (WAO) taskforce. *Allergy*. 2007;62(3):317-24.
  53. Clark J, Schall R. Assessment of combined symptom and medication scores for rhinoconjunctivitis immunotherapy clinical trials. *Allergy*. 2007;62(9):1023-8.
  54. Casale TB, Romero FA, Spierings EL. Intranasal noninhaled carbon dioxide for the symptomatic treatment of seasonal allergic rhinitis. *J Allergy Clin Immunol*. 2008;121(1):105-9.
  55. Molgaard E, Thomsen SF, Lund T, Pedersen L, Nolte H, Backer V. Differences between allergic and nonallergic rhinitis in a large sample of adolescents and adults. *Allergy*. 2007;62(9):1033-7.
  56. Greiner AN, Meltzer EO. Pharmacologic rationale for treating allergic and nonallergic rhinitis. *J Allergy Clin Immunol*. 2006;118(5):985-98.
  57. Linneberg A, Jorgensen T, Nielsen NH, Madsen F, Frolund L, Dirksen A. The prevalence of skin-test-positive allergic rhinitis in Danish adults: two cross-sectional surveys 8 years apart. The Copenhagen Allergy Study. *Allergy*. 2000;55(8):767-72.
  58. Terreehorst I, Oosting AJ, Tempels-Pavlica Z, de Monchy JG, Bruijnzeel-Koomen CA, Hak E, van Wijk RG. Prevalence and severity of allergic rhinitis in house dust mite-allergic patients with bronchial asthma or atopic dermatitis. *Clin Exp Allergy*. 2002;32(8):1160-5.
  59. Georgy V, Fahim HI, El Gaafary M, Walters S. Prevalence and socioeconomic associations of asthma and allergic rhinitis in Cairo, Egypt. *Eur Respir J*. 2006;28(4):756-62.
  60. Kuyucu S, Saraclar Y, Tuncer A, Geyik PO, Adalioğlu G, Akpinarlı A, Sekerel BE, Sümbülo lu V. Epidemiologic characteristics of rhinitis in Turkish children: the International Study of Asthma and Allergies in Childhood (ISAAC) phase 2. *Pediatr Allergy Immunol*. 2006;17(4):269-77.
  61. Antonicelli L, Micucci C, Voltolini S, Feliziani V, Senna GE, Di Blasi P, Visonà G, De Marco R, Bonifazi F. Allergic rhinitis and asthma comorbidity: ARIA classification of rhinitis does not correlate with the prevalence of asthma. *Clin Exp Allergy*. 2007;37(6):954-60.
  62. Antonicelli L, Micucci C, Voltolini S, Senna GE, Di Blasi P, Visonà G, De Marco R, Bonifazi F. Relationship between ARIA classification and drug treatment in allergic rhinitis and asthma. *Allergy*. 2007;62(9):1064-70.
  63. Stelmach R, do Patrocinio TNM, Ribeiro M, Cukier A. Effect of treating allergic rhinitis with corticosteroids in patients with mild-to-moderate persistent asthma. *Chest*. 2005;128(5):3140-7.
  64. Camargos P, Ibiapina C, Lasmar L, Cruz AA. Obtaining concomitant control of allergic rhinitis and asthma with a nasally inhaled corticosteroid. *Allergy*. 2007;62(3):310-6.
  65. Jacobsen L, Niggemann B, Dreborg S, Ferdousi HA, Halken S, Host A, Koivikko A, Norberg LA, Valovirta E, Wahn U, Möller C; (The PAT investigator group). Specific immunotherapy has long-term preventive effect of seasonal and perennial asthma: 10-year follow-up on the PAT study. *Allergy*. 2007;62(8):943-8.
  66. Civelek E, Soyer OU, Gemicioglu B, Sekerel BE. Turkish physicians' perception of allergic rhinitis and its impact on asthma. *Allergy*. 2006;61(12):1454-8.
  67. Demoly P, Concas V, Urbinelli R, Allaert F. Evaluation de l'influence des recommandations OMS-ARIA sur la prise en charge de la rhinite allergique en pratique de ville en France. *Enquête ERNANI. Rev Fr Allergol Immunol Clin*. 2006;46:626-32.
  68. Pereira C, Valero A, Loureiro C, Davila I, Martinez-Cocera, Murio C, Rico P, Palomino R. Iberian study of aeroallergens sensitisation in allergic rhinitis. *Allerg Immunol*. 2006;38:186-194.

69. Workshop Report, Global Strategy for Asthma Management and Prevention (GINA). Initiative World Health Organization, WHO. Updated November 2006. <http://www.ginasthma.com/>
70. Alergológica 1995. Factores Epidemiológicos Clínicos y Socioeconómicos de las enfermedades alérgicas en España en 1995. Sociedad Española de Alergología e Inmunología Clínica y Alergia e Inmunología Clínica (Eds). Madrid, 1995.
71. Alergológica 2005. Factores Epidemiológicos Clínicos y Socioeconómicos de las enfermedades alérgicas en España en 2005. Sociedad Española de Alergología e Inmunología Clínica y Alergia e Inmunología Clínica. (Eds). Madrid 2005.
72. Navarro A, Valero A, Juliá B, Quirce S. Coexistence of asthma and allergic rhinitis in adult patients attending allergy clinics: ONEAIR Study. *J Invest Allergol Clin Immunol*. 2008 (In press).
73. Castillo JA; Mullol J. Comorbilidad de rinitis y asma en España. *Arch Bronconeumol*. 2008 (In press).
74. Valero A, Alonso J, Antépara I, Baró E, Colás C, del Cuvillo A, Ferrer M, Herdman M, Marti-Guadaño E, Monclús L, Mullol J, Navarro-Pulido AM, Navas C, Sastre J and the ESPRINT Group of Investigators. Development and validation of a new Spanish instrument to measure health-related quality of life in patients with allergic rhinitis: The ESPRINT Questionnaire. *Val Health*. 2007;10:466-477.
75. Valero A, Alonso J, Antépara I, Baró E, Colás C, del Cuvillo A, Ferrer M, Herdman M, Marti-Guadaño E, Monclús L, Navarro-Pulido AM, Sastre J, Izquierdo I, Mullol J. Health-related quality of life in allergic rhinitis: comparing the short form ESPRINT-15 and MiniRQLQ questionnaires. *Allergy*. 2007;62(12):1372-8.
76. Togias A. Mechanistic features of allergic rhinitis. *J Allergy Clin Immunol*. 2000; 105:S559-604.
77. Juniper EF, Guyatt GH. Development and testing of a new measure of health status for clinical trials in rhinoconjunctivitis. *Clin Exp Allergy*. 1991;21:77-83.
78. Juniper EF, Guyatt GH, Dolovich J. Assessment of quality of life in adolescents with allergic rhinoconjunctivitis: development and testing of a questionnaire for clinical trials. *J Allergy Clin Immunol*. 1994; 93(2):413-23.
79. Juniper EF, Howland WC, Roberts NB. Measuring quality of life in children with rhinoconjunctivitis. *J Allergy Clin Immunol*. 1998;101:163-70.
80. Juniper EF, Thompson AK, Ferrie PJ, Roberts JN. Development and validation of the mini rhinoconjunctivitis quality of life questionnaire. *Clin Exp Allergy*. 2000;30(1):132-40.
81. Thomson AK, Juniper E, Meltzer EO. Quality of life in patients with allergic rhinitis. *Ann Allergy Asthma Immunol*. 2000;85:338-48.
82. Vilagut G, Ferrer M, Rajmil L, Rebollo P, Permanyer-Miralda G, Quintana JM, Santed R, Valderas JM, Ribera A, Domingo-Salvany A, Alonso J. The Spanish version of the Short Form 36 Health Survey: a decade of experience and new developments. *Gac Sanit*. 2005;19(2):135-50.
83. Bachert C, Bousquet J, Canonica GW, Durham SR, Klimek L, Mullol J, Van Cauwenberge PB, Van Hammée G; XPERT Study Group. Levocetirizine improves quality of life and reduces costs in long-term management of persistent allergic rhinitis. *J Allergy Clin Immunol*. 2004;114:838-44.
84. Schramm B, Ehlken B, Smala A, Quednau K, Berger K, Nowak D. Cost of illness of atopic asthma and seasonal allergic rhinitis in Germany: 1-yr retrospective study. *Eur Respir J*. 2003;21:116-122.
85. Halpern MT, Schmier JK, Richner R, Guo C, Togias A. Allergic rhinitis: a potential cause of increased asthma medication use, costs, and morbidity. *J Asthma*. 2004;41:117-126.

#### ■ Joaquim Mullol

Unitat de Rinologia i Clínica de l'Olfacte  
Servei d'ORL, Hospital Clínic  
C/ Villarroel 170  
08036 Barcelona  
Catalonia, Spain  
E-mail: [jmullol@clinic.ub.es](mailto:jmullol@clinic.ub.es)